

Anti-GAP43 Antibody

Our Anti-GAP43 primary antibody from PhosphoSolutions is chicken polyclonal. It detects bovine, hors Catalog # AN1410

Specification

Anti-GAP43 Antibody - Product Information

Application	WB, IHC
Primary Accession	<u>P07936</u>
Reactivity	Bovine, Chicken, Drosophila
Host	Chicken
Clonality	Polyclonal
Isotype	IgY
Calculated MW	23603

Anti-GAP43 Antibody - Additional Information

Gene ID

29423

Other Names

Axonal membrane protein GAP 43 antibody, Axonal membrane protein GAP-43 antibody, B 50 antibody, Calmodulin binding protein P 57 antibody, F1 antibody, GAP 43 antibody, GAP-43 antibody, GAP43 antibody, Growth Associated Protein 43 antibody, Growth-associated protein 43 antibody, Nerve Growth Related Peptide antibody, Nerve growth related peptide GAP43 antibody, NEUM_HUMAN antibody, Neural phosphoprotein B 50 antibody, Neural phosphoprotein B-50 antibody, Neuromodulin antibody, Neuron growth associated protein 43 antibody, PP46 antibody, Protein F1 antibody, QtrA-11580 antibody, QtrA-13071 antibody

Target/Specificity

GAP-43 is thought to have an important role in development and plasticity because it is expressed at high levels in neuronal growth cones during development and during axonal regeneration (Benowitz and Routtenberg, 1997). There is also evidence from knockout animals that GAP-43 serves to amplify pathfinding signals from the growth cone (Strittmatter et al., 1995). GAP-43 is thought to mediate at least some of these effects via interaction with actin. Importantly, phosphorylation at Ser-41 by protein kinase C modulates the interaction of GAP-43 with actin (He et al., 1997) and may also affect neurotransmitter release during forms of plasticity like LTP (Hulo et al., 2002).

Dilution WB~~1:1000 IHC~~1:100~500

Format Total IgY fraction

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-GAP43 Antibody is for research use only and not for use in diagnostic or therapeutic



procedures.

Shipping Blue Ice

Anti-GAP43 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

Anti-GAP43 Antibody - Images



Western blot of rat cortical lysate showing specific immunolabeling of the \sim 43 kDa GAP43 protein.



Immunolabeling of mixed neuron and glial cultures with anti-GAP43 (cat. 875-GAP43, 1:2000, green), alpha II spectrin (cat. 99-A2SR, 1:1000, red) and nuclear staining with DAPI (blue). The anti-GAP43 labels protein expressed in the axonal membrane and synapses of neuronal cells. The Anti- α II-Spectrin labels the submembraneous cytoskeleton of axons and dendrites.



Immunofluorescence of rat neuron/glial culture labeling GAP43 protein (cat. 875-GAP43, 1:2000, green) expressed in the axonal membrane and synapses of neuronal cells. DAPI (blue) was used for nuclear staining.

Anti-GAP43 Antibody - Background

GAP-43 is thought to have an important role in development and plasticity because it is expressed at high levels in neuronal growth cones during development and during axonal regeneration (Benowitz and Routtenberg, 1997). There is also evidence from knockout animals that GAP-43 serves to amplify pathfinding signals from the growth cone (Strittmatter et al., 1995). GAP-43 is thought to mediate at least some of these effects via interaction with actin. Importantly, phosphorylation at Ser-41 by protein kinase C modulates the interaction of GAP-43 with actin (He et al., 1997) and may also affect neurotransmitter release during forms of plasticity like LTP (Hulo et al., 2002).