

Anti-NMDA NR2C Subunit (Ser1244) Antibody

Our Anti-NMDA NR2C Subunit (Ser1244) rabbit polyclonal phosphospecific primary antibody from Phospho Catalog # AN1495

Specification

Anti-NMDA NR2C Subunit (Ser1244) Antibody - Product Information

Primary Accession Host Clonality	<u>000961</u> Rabbit Polyclonal IgG
Calculated MW	135271

Anti-NMDA NR2C Subunit (Ser1244) Antibody - Additional Information

Gene ID

24411

Other Names

GluN2C antibody, Glutamate [NMDA] receptor subunit epsilon-3 antibody, Glutamate Receptor Ionotropic N Methyl D Aspartate 2C antibody, Glutamate receptor ionotropic NMDA 2C antibody, Glutamate receptor ionotropic N methyl D aspartate 2C antibody, Grin2c antibody, Grin 2c antibody, GRIN2C antibody, GRIN 2C antibody, N Methly D Aspartate Receptor Channel Subunit Epsilon 3 antibody, N Methyl D Aspartate Receptor Subtype 2C antibody, N methyl D aspartate receptor subunit 2C antibody, N-methyl D-aspartate receptor subtype 2C antibody, NMDAR2C antibody, NMDE3 HUMAN antibody, NR2C antibody, OTTHUMP00000041930 antibody

Target/Specificity

The ion channels activated by glutamate that are sensitive to N-methyl-Daspartate (NMDA) are designated NMDA receptors (NMDAR). The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). The NMDA receptor is also one of the principal molecular targets for alcohol in the CNS (Lovinger et al., 1989; Alvestad et al., 2003; Snell et al., 1996). The NMDAR is also potentiated by protein phosphorylation (Lu et al., 1999). The NR2C subunit of the receptor is thought to influence the NMDAR conductance level (Ebralidze et al., 1996). Phosphorylation of Ser-1244 has been shown to regulate NMDA receptor channel function (Chen et al., 2006).

Format

Antigen Affinity Purified from Pooled Serum

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-NMDA NR2C Subunit (Ser1244) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice



Anti-NMDA NR2C Subunit (Ser1244) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-NMDA NR2C Subunit (Ser1244) Antibody - Images



Western blot of rat cerebellar lysate showing specific immunolabeling of the \sim 140 kDa NR2C subunit of the NMDA receptor phosphorylated at Ser1244 in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is blocked by preadsorption of the phosphopeptide used as the antigen, but not by the corresponding non-phosphopeptide (not shown).

Anti-NMDA NR2C Subunit (Ser1244) Antibody - Background

The ion channels activated by glutamate that are sensitive to N-methyl-Daspartate (NMDA) are designated NMDA receptors (NMDAR). The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). The NMDA receptor is also one of the principal molecular targets for alcohol in the CNS (Lovinger et al., 1989; Alvestad et al., 2003; Snell et al., 1996). The NMDAR is also potentiated by protein phosphorylation (Lu et al., 1999). The NR2C subunit of the receptor is thought to influence the NMDAR conductance level (Ebralidze et al., 1996). Phosphorylation of Ser-1244 has been shown to regulate NMDA receptor channel function (Chen et al., 2006).