

Anti-Troponin I (cardiac) Ser23/24 Antibody

Our Anti-Troponin I (cardiac) Ser23/24 rabbit polyclonal phosphospecific primary antibody from Phosp
Catalog # AN1589

Specification**Anti-Troponin I (cardiac) Ser23/24 Antibody - Product Information**

Primary Accession	P48787
Reactivity	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	24259

Anti-Troponin I (cardiac) Ser23/24 Antibody - Additional Information

Gene ID **21954**

Other Names

cardiac muscle antibody, Cardiac troponin I antibody, Cardiac Troponin I antibody, cardiomyopathy dilated 2A (autosomal recessive) antibody, Cardiomyopathy familial hypertrophic 7 included antibody, CMD1FF antibody, CMD2A antibody, CMH7 antibody, cTnI antibody, Familial hypertrophic cardiomyopathy 7 antibody, MGC116817 antibody, RCM1 antibody, Tn1 antibody, Tni antibody, TNN I3 antibody, TNNC 1 antibody, TNNC1 antibody, TNNI3 antibody, TNNI3_HUMAN antibody, Troponin I antibody, Troponin I cardiac antibody, Troponin I cardiac muscle antibody, Troponin I cardiac muscle isoform antibody, Troponin I type 3 cardiac antibody, troponin I cardiac 3 antibody, TroponinI antibody, Troponin I type 3 (cardiac) antibody

Target/Specificity

Troponin I (cTnI) is 1 of 3 subunits, along with troponin C (TnC) and troponin T (TnT) of troponin complex found in cardiac muscle. cTnI binds to actin in thin myofilaments to hold the troponin-tropomyosin complex in place. Phosphorylation of cardiac isoform of TnI at serines 22,23 in the unique amino-terminal end molecule decreases the calcium sensitivity of the sarcomere, promotes calcium dissociation from troponin C and by extension enhances rates of cross-bridge cycling and diastolic relaxation (Noland, Jr. et al., 1995; Noland et al., 1989). In addition, studies using reconstituted fibers and mutational analysis have shown that PKC phosphorylation of TnI (largely at Ser-43) inhibits the actin-cross bridge reaction and reduces the Ca⁺⁺ dependent actomyosin ATPase rate as well as the calcium sensitivity of force generation (Noland, Jr. and Kuo, 1991). Phosphorylation at Thr-144 (mediated by several PKC isoforms) reduces maximal tension development and cross-bridge cycling rates (Sumandea et al., 2008). Importantly, changes in the phosphorylation at each of these sites have been shown to be stage-specific with regard to cardiac disease progression (Walker et al., 2010).

Format

Antigen Affinity Purified from Pooled Serum

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Troponin I (cardiac) Ser23/24 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Shipping
 Blue Ice

Anti-Troponin I (cardiac) Ser23/24 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Troponin I (cardiac) Ser23/24 Antibody - Images



Western blot of mouse heart lysate showing specific immunolabeling of the ~25 kDa cardiac troponin I protein phosphorylated at Ser23/24 in the second lane (-). Phosphospecificity is shown in the first lane (+) where immunolabeling is blocked by preadsorption with the phosphopeptide used as antigen, but not by the corresponding non-phosphopeptide (not shown).



Western blots showing specific labeling of S23/S24 TnI, cat. p2010-2324 (upper) and TnI; cat. 2010-TnI (lower) in protein lysates of mouse cardiomyocytes. Image from publication CC-BY-4.0. PMID: 36424917

Anti-Troponin I (cardiac) Ser23/24 Antibody - Background

Troponin I (cTnI) is 1 of 3 subunits, along with troponin C (TnC) and troponin T (TnT) of troponin complex found in cardiac muscle. cTnI binds to actin in thin myofilaments to hold the troponin-tropomyosin complex in place. Phosphorylation of cardiac isoform of TnI at serines 22,23 in the unique amino-terminal end molecule decreases the calcium sensitivity of the sarcomere, promotes calcium dissociation from troponin C and by extension enhances rates of cross-bridge cycling and diastolic relaxation (Noland, Jr. et al., 1995; Noland et al., 1989). In addition, studies using reconstituted fibers and mutational analysis have shown that PKC phosphorylation of TnI (largely at Ser-43) inhibits the actin-cross bridge reaction and reduces the Ca⁺⁺ dependent actomyosin ATPase rate as well as the calcium sensitivity of force generation (Noland, Jr. and Kuo, 1991). Phosphorylation at Thr-144 (mediated by several PKC isoforms) reduces maximal tension development and cross-bridge cycling rates (Sumandea et al., 2008). Importantly, changes in the phosphorylation at each of these sites have been shown to be stage-specific with regard to cardiac disease progression (Walker et al., 2010).