

Anti-Akt (Ser-473), Phosphospecific Antibody

Catalog # AN1625

Specification

Anti-Akt (Ser-473), Phosphospecific Antibody - Product Information

Application	WB
Primary Accession	<u>P31749</u>
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	lgG1
Calculated MW	55686

Anti-Akt (Ser-473), Phosphospecific Antibody - Additional Information

Gene ID Other Names PKBalpha, PKB, AKT 207

Target/Specificity

Akt (PKB, Rac kinase) is a 60kDa ser/thr kinase critical for controlling diverse cellular functions, including glucose metabolism, gene transcription, cell proliferation, and apoptosis. Akt phosphorylates a number of substrates including MBP, glycogen synthetase, PKA RII subunit, and histone H1. Akt is activated in response to insulin and growth factors in a PI3-kinase dependent manner. Activation of PI3-Kinase generates phosphatidylinositol 3,4-bisphosphate, which induces membrane translocation of Akt coincident with its phosphorylation at Thr-308 and Ser-473. Upon activation, Akt associates with members of the PKC family of kinases, such as PKCδ and PKCζ. Ceramide-activated PKCζ leads to phosphorylation of Thr-34 within the pleckstrin homology domain of Akt. This phosphorylation inhibits PIP3 binding to Akt preventing activation of the kinase and may lead to cermide-induced cell death.

Dilution WB~~1:1000

Format Protein G Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Akt (Ser-473), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice



Anti-Akt (Ser-473), Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Akt (Ser-473), Phosphospecific Antibody - Images



Western blot analysis of A431 cells untreated (lanes 1 & 3) or treated with 100 ng/ml EGF for 60 min. (lanes 2 & 4). The blots were probed with monoclonal anti-phospho-Akt (Ser-473) (lanes 1 & 2) and monoclonal anti-Akt1 (N-terminal region) (lanes 3 & 4).

Anti-Akt (Ser-473), Phosphospecific Antibody - Background

Akt (PKB, Rac kinase) is a 60kDa ser/thr kinase critical for controlling diverse cellular functions, including glucose metabolism, gene transcription, cell proliferation, and apoptosis. Akt phosphorylates a number of substrates including MBP, glycogen synthetase, PKA RII subunit, and histone H1. Akt is activated in response to insulin and growth factors in a PI3-kinase dependent manner. Activation of PI3-Kinase generates phosphatidylinositol 3,4-bisphosphate, which induces membrane translocation of Akt coincident with its phosphorylation at Thr-308 and Ser-473. Upon activation, Akt associates with members of the PKC family of kinases, such as PKCδ and PKCζ. Ceramide-activated PKCζ leads to phosphorylation of Thr-34 within the pleckstrin homology domain of Akt. This phosphorylation inhibits PIP3 binding to Akt preventing activation of the kinase and may lead to cermide-induced cell death.