

Anti-Akt (Thr-34), Phosphospecific Antibody
Catalog # AN1626**Specification**

Anti-Akt (Thr-34), Phosphospecific Antibody - Product Information

| | |
|-------------------|--------------------------|
| Primary Accession | P31749 |
| Reactivity | Bovine |
| Host | Rabbit |
| Clonality | Rabbit Polyclonal |
| Isotype | IgG |
| Calculated MW | 55686 |

Anti-Akt (Thr-34), Phosphospecific Antibody - Additional Information

| | |
|--------------------|------------|
| Gene ID | 207 |
| Other Names | |
| PKBalpha, PKB, AKT | |

Target/Specificity

Akt (PKB, Rac kinase) is a 60kDa ser/thr kinase critical for controlling diverse cellular functions, including glucose metabolism, gene transcription, cell proliferation, and apoptosis. Akt phosphorylates a number of substrates including MBP, glycogen synthetase, PKA RII subunit, and histone H1. Akt is activated in response to insulin and growth factors in a PI3-kinase dependent manner. Activation of PI3-Kinase generates phosphatidylinositol 3,4-bisphosphate, which induces membrane translocation of Akt coincident with its phosphorylation at Thr-308 and Ser-473. Upon activation, Akt associates with members of the PKC family of kinases, such as PKC δ and PKC ζ . Ceramide-activated PKC ζ leads to phosphorylation of Thr-34 within the pleckstrin homology domain of Akt. This phosphorylation inhibits PIP3 binding to Akt preventing activation of the kinase and may lead to ceramide-induced cell death.

Format

Antigen Affinity Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Akt (Thr-34), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-Akt (Thr-34), Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Akt (Thr-34), Phosphospecific Antibody - Images



Western blot analysis of A431 cells, serum starved overnight (20 µg/lanes 1 & 3) and calyculin A (100 nM) treated for 30 minutes (20 µg/lanes 2 & 4). The blot was probed with anti-Akt (Thr-34) (lanes 1 & 2) or anti-Akt1 (N-terminal region) (lanes 3 & 4).

Anti-Akt (Thr-34), Phosphospecific Antibody - Background

Akt (PKB, Rac kinase) is a 60kDa ser/thr kinase critical for controlling diverse cellular functions, including glucose metabolism, gene transcription, cell proliferation, and apoptosis. Akt phosphorylates a number of substrates including MBP, glycogen synthetase, PKA RII subunit, and histone H1. Akt is activated in response to insulin and growth factors in a PI3-kinase dependent manner. Activation of PI3-Kinase generates phosphatidylinositol 3,4-bisphosphate, which induces membrane translocation of Akt coincident with its phosphorylation at Thr-308 and Ser-473. Upon activation, Akt associates with members of the PKC family of kinases, such as PKCδ and PKCζ. Ceramide-activated PKCζ leads to phosphorylation of Thr-34 within the pleckstrin homology domain of Akt. This phosphorylation inhibits PIP3 binding to Akt preventing activation of the kinase and may lead to ceramide-induced cell death.