

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody Catalog # AN1748

Specification

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody - Product Information

Primary Accession	
Reactivity	
Host	
Clonality	
Isotype	
Calculated MW	

<u>099704</u> Bovine Rabbit Rabbit Polyclonal IgG 52392

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody - Additional Information

Gene ID Other Names p62DOK

1796

Target/Specificity

Doks are a family of adaptor proteins that include six Dok proteins (Dok1 to Dok6), which have an N-terminal pleckstrin homology domain, a central phosphotyrosine binding domain, and a C-terminal region containing multiple tyrosine residues. When phosphorylated, these tyrosines can serve as docking sites for SH2 domain-containing proteins. Dok1 (p62dok) has been shown to bind Ras-GAP, Nck, and Csk. Several tyrosine phosphorylation sites have been identified for Dok1. One site, Tyr-362 (Tyr-361 mouse), is phosphorylated by c-Abl, is required for Nck binding, and may be critical for filopodia formation during fibroblast spreading on fibronectin. Alternatively, Dok1 activity is also regulated by serine phosphorylation. IkB Kinase β phosphorylates several serine sites including Ser-450 in vitro, and TNF α , IL-1, and radiation treatment lead to phosphorylation of Ser-443, Ser-446, and Ser-450 in vivo. Phosphorylation of these serine sites may be required for Dok-mediated inhibition of MAPK signaling and stimulation of cell motility.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides



- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody - Images



Western blot image of Jurkat cells stimulated with calyculin A (100 nM, 30 min) (lanes 1-6) followed by lambda phosphatase (lanes 2 & 6) or alkaline phosphatase (lane 4) treatment. The blots were probed with anti-Dok1 (lanes 1 & 2), anti-Dok1 (Tyr-362) (lanes 3 & 4), and anti-Dok1 (Ser-450) (lanes 5 & 6).

Anti-Dok1 (Tyr-362) [Dok2 (Tyr-337)], Phosphospecific Antibody - Background

Doks are a family of adaptor proteins that include six Dok proteins (Dok1 to Dok6), which have an N-terminal pleckstrin homology domain, a central phosphotyrosine binding domain, and a C-terminal region containing multiple tyrosine residues. When phosphorylated, these tyrosines can serve as docking sites for SH2 domain-containing proteins. Dok1 (p62dok) has been shown to bind Ras-GAP, Nck, and Csk. Several tyrosine phosphorylation sites have been identified for Dok1. One site, Tyr-362 (Tyr-361 mouse), is phosphorylated by c-Abl, is required for Nck binding, and may be critical for filopodia formation during fibroblast spreading on fibronectin. Alternatively, Dok1 activity is also regulated by serine phosphorylation. IkB Kinase β phosphorylates several serine sites including Ser-450 in vitro, and TNF α , IL-1, and radiation treatment lead to phosphorylation of Ser-443, Ser-446, and Ser-450 in vivo. Phosphorylation of these serine sites may be required for Dok-mediated inhibition of MAPK signaling and stimulation of cell motility.