

Anti-eEF2K (Ser-398), Phosphospecific Antibody
Catalog # AN1765**Specification****Anti-eEF2K (Ser-398), Phosphospecific Antibody - Product Information**

Primary Accession	O00418
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	82144

Anti-eEF2K (Ser-398), Phosphospecific Antibody - Additional InformationGene ID **29904****Other Names**

eEF-2, eEF-2K, CaMK-III, eukaryotic elongation factor

Target/Specificity

Eukaryotic elongation factor 2 (eEF2) catalyzes the translocation of peptidyl-tRNA from the A site to the P site on the ribosome. eEF2 kinase (eEF2K) phosphorylates and inactivates eEF2, resulting in the inhibition of peptide-chain elongation. eEF2K is normally dependent on Ca²⁺ ions and calmodulin, and can be activated by PKA in response to elevated cAMP levels during cell stress- or starvation-related conditions. Regulation of eEF2K occurs through phosphorylation at multiple sites. Ser-78 phosphorylation is required for calmodulin binding and eEF2K activity, while phosphorylation of Ser-500 is required for Ca²⁺/calmodulin-independent kinase activity. Thr-348 is an autophosphorylation site that is required for kinase activity. Inhibitory phosphorylation may also regulate eEF2K, since phosphorylation at Ser-359 by SAPK4/p38 δ causes inactivation of eEF2K. Thus, multisite phospho-regulation of eEF2K may be important for proper control of eEF2K activity and protein translation.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-eEF2K (Ser-398), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice


Anti-eEF2K (Ser-398), Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-eEF2K (Ser-398), Phosphospecific Antibody - Images



Western blot image of human Jurkat cells treated with calyculin A (100 nM) for 30 min. (lanes 1-6), then some lanes of the blot were treated with lambda phosphatase (lanes 2, 4 & 6). The blot was probed with rabbit polyclonals anti-eEF2K (C-terminus) (lanes 1 & 2), anti-eEF2K (Ser-445) phospho-specific (lanes 3 & 4), and anti-eEF2K (Ser-398) phospho-specific (lanes 5 & 6).

Anti-eEF2K (Ser-398), Phosphospecific Antibody - Background

Eukaryotic elongation factor 2 (eEF2) catalyzes the translocation of peptidyl-tRNA from the A site to the P site on the ribosome. eEF2 kinase (eEF2K) phosphorylates and inactivates eEF2, resulting in the inhibition of peptide-chain elongation. eEF2K is normally dependent on Ca^{2+} ions and calmodulin, and can be activated by PKA in response to elevated cAMP levels during cell stress- or starvation-related conditions. Regulation of eEF2K occurs through phosphorylation at multiple sites. Ser-78 phosphorylation is required for calmodulin binding and eEF2K activity, while phosphorylation of Ser-500 is required for Ca^{2+} /calmodulin-independent kinase activity. Thr-348 is an autophosphorylation site that is required for kinase activity. Inhibitory phosphorylation may also regulate eEF2K, since phosphorylation at Ser-359 by SAPK4/p38 δ causes inactivation of eEF2K. Thus, multisite phospho-regulation of eEF2K may be important for proper control of eEF2K activity and protein translation.