

Anti-Plexin D1 (C-terminal region) Antibody
Catalog # AN1917**Specification****Anti-Plexin D1 (C-terminal region) Antibody - Product Information**

Primary Accession	Q9Y4D7
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	212007

Anti-Plexin D1 (C-terminal region) Antibody - Additional Information

Gene ID	23129
Other Names	
PLXND1, Sema3E	

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Plexin D1 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-Plexin D1 (C-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Plexin D1 (C-terminal region) Antibody - Images



Western blot analysis of Plexin D1 expression in human endothelial cells (HUVEC) (lanes 1-6). The blots were probed with rabbit polyclonals anti-Plexin D1 (Sema domain) (lanes 1 & 2), anti-Plexin D1 (Cytoplasmic domain) (lanes 3 & 4), and anti-Plexin D1 (a.a. 1635-1647) (lanes 5 & 6). Each antibody was used in the presence of their respective blocking peptide (lanes 2, 4 & 6).

Anti-Plexin D1 (C-terminal region) Antibody - Background

Plexins are a family of large integral membrane proteins that complex with neuropilins to form semaphorin co-receptors. The extracellular region of plexins contains a semaphorin domain, multiple glycine-rich motifs, and MET-related sequences. The cytoplasmic region contains a Sex/Plexin domain and putative tyrosine phosphorylation sites that mediate signal transduction after activation. Plexin D1 is a type I transmembrane protein that may be expressed as multiple isoforms in many cell types, including neurons and endothelial cells. Semaphorin 3E (Sema-3E) and semaphorin 4A can bind Plexin D1, and ligand binding leads to phosphorylation of Tyr-1642 and activation of Plexin D1. Sema-3E and its receptor are important for angiogenesis that occurs during blood vessel development and repair. In cancers, Sema-3E and Plexin D1 are expressed in tumor tissues and cancer cells, and the interaction of these molecules may promote cancer cell migration and metastatic spreading. Thus, Plexin D1, and its ligand Sema-3E, may be important regulators of angiogenesis and metastasis.