

Anti-PYK2 (C-terminal region) Antibody
Catalog # AN1928**Specification****Anti-PYK2 (C-terminal region) Antibody - Product Information**

Application	WB
Primary Accession	Q14289
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Calculated MW	115875

Anti-PYK2 (C-terminal region) Antibody - Additional Information

Gene ID	2185
Other Names	
CAKbeta, FAK2	

Dilution

WB~~1:1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-PYK2 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Shipping


Blue Ice

Anti-PYK2 (C-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PYK2 (C-terminal region) Antibody - Images



Western blot analysis of mouse brain. The blot was probed with anti-PYK2 (C-terminal region) antibody at 1:250 (lane 1), 1:500 (lane 2), 1:1000 (lane 3), and 1:2000 (lane 4).

Anti-PYK2 (C-terminal region) Antibody - Background

Protein tyrosine kinase, Pyk2 (CAKb, RAFTK, CADTK), is a nonreceptor tyrosine kinase structurally related to focal adhesion kinase (FAK). Pyk2 is predominantly expressed in cells derived from hematopoietic lineages and in the central nervous system. Pyk2 is one of the signaling mediators for the G-protein-coupled receptors and MAP kinase signaling pathway. It plays an important role in cell spreading and migration of various cell types. In T-cells, Pyk2 is tyrosine phosphorylated and activated upon ligation of TCR. Phosphorylation of Tyr-402 is required for the phosphorylation of other tyrosines in Pyk2 and provides a binding site for Fyn SH2 during T-cell activation.