

# Anti-Semaphorin-3A (C-terminal) Antibody

Catalog # AN1944

# **Specification**

### Anti-Semaphorin-3A (C-terminal) Antibody - Product Information

Primary Accession

Host

O14563

Rabbit

Clonality Rabbit Polyclonal

Isotype IgG
Calculated MW 88889

# Anti-Semaphorin-3A (C-terminal) Antibody - Additional Information

Gene ID **10371** 

**Other Names** 

Sema3A, Semaphorin III

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

Anti-Semaphorin-3A (C-terminal) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping** 

Blue Ice


## Anti-Semaphorin-3A (C-terminal) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Anti-Semaphorin-3A (C-terminal) Antibody - Images





Western blot image of human A431 cells and (lane 1) and human recombinant Sema3A/Fc chimera (95/125 kDa) (lane 2). The blots were probed with rabbit polyclonal anti-Semaphorin 3A (SP1241) at a dilution of 1:1000.

# Anti-Semaphorin-3A (C-terminal) Antibody - Background

One family of inhibitory axon guidance molecules is the semaphorins. The semaphorins include secreted, transmembrane, and GPI-anchored extracellular molecules that are involved in regulating axon guidance by inhibiting axons from growing toward incorrect targets. Semaphorin 3A (Sema3A) may play a particularly interesting role in limiting axon regeneration since it is expressed in meningeal fibroblasts that invade the injured spinal cord and surround the glial scar. In addition, the Sema3A co-receptors, Neuropilin-1 and Plexin-A1, are expressed on axons that regenerate up to the injured region, but do not cross this Sema3A-containing region. Thus, Sema3A and its co-receptors may have important roles in regulating axon guidance during neuronal development and after neuronal injury.