

Anti-Semaphorin-6A (C-terminus) Antibody
Catalog # AN1950**Specification****Anti-Semaphorin-6A (C-terminus) Antibody - Product Information**

Application	WB, IHC
Primary Accession	Q9H2E6
Reactivity	Bovine, Chicken
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	114369

Anti-Semaphorin-6A (C-terminus) Antibody - Additional Information

Gene ID 57556

Dilution

WB~~1:1000

IHC~~1:100~500

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Semaphorin-6A (C-terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Shipping


Blue Ice

Anti-Semaphorin-6A (C-terminus) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Semaphorin-6A (C-terminus) Antibody - Images



Western blot of Sema-6A in human colorectal cancer (CRC) cells treated with control (shC) or Sema-6A (shSema6A) shRNAs. Sema-6A was immunoprecipitated from each of the CRC lysates using Sema-6A (a.a. 772-787) or Sema-6A (C-terminus) antibody, then the blotted immunoprecipitations were probed with Sema-6A antibody. (Images provided by Dr. Luca Tamagnone from the IRCC, Univ. of Torino, Italy).



Immunocytochemical labeling of Sema-6A in COS7 cells that were mock transfected (top images) or Sema-6A transfected (bottom images). The cells were labeled with anti-Sema-6A (a.a. 772-787) (Left top and bottom image) or anti-Sema-6A (C-terminus) (Right top and bottom image). The antibodies were detected using anti-rabbit fluorescent secondary antibody. (Images provided by Dr. Luca Tamagnone from the IRCC, University of Torino, Italy).

Anti-Semaphorin-6A (C-terminus) Antibody - Background

The Semaphorin family of axon guidance molecules includes secreted, transmembrane, and GPI-anchored extracellular molecules that have been implicated in neuron development, vascular disease, and tumor progression. There are eight classes of semaphorin genes, all of which are characterized by a conserved 500 amino acid, cystine-rich Sema domain. Semaphorin 6A (Sema-6A) is a single-pass type I membrane protein that contains a plexin-semaphorin-integrin (PSI) domain and a Sema domain. Sema-6A is active as a homodimer or oligomer. The Sema-6A homodimer interacts with a Plexin A2 homodimer. Sema-6a promotes reorganization of the actin cytoskeleton to regulate normal granule cell migration in the developing cerebellum and to control axon guidance in the developing central nervous system. In addition, Sema-6a has been implicated in oligodendrocyte myelination and may alter VEGF signaling during angiogenesis.