

Anti-SHP1 (C-terminal region) Antibody
Catalog # AN1953**Specification**

Anti-SHP1 (C-terminal region) Antibody - Product Information

Application	WB, IHC
Primary Accession	P29350
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Calculated MW	67561

Anti-SHP1 (C-terminal region) Antibody - Additional Information

Gene ID	5777
Other Names	
PTP1C, SHPTP1	

Dilution

WB~~1:1000
IHC~~1:100~500

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-SHP1 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-SHP1 (C-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SHP1 (C-terminal region) Antibody - Images



Western blot analysis of human Jurkat cells treated with pervanadate (1 mM) for 30 min. The blot was exposed to lambda phosphatase (lanes 2 & 4) then probed with anti-SHP1 (C-terminal) antibody (lanes 1 & 2) or anti-SHP1 (Ser-591) antibody (lanes 3-6). The SHP1 (Ser-591) antibody was used in the presence of phospho-SHP1 (Ser-591) peptide (lane 5) or a non-specific phospho-serine peptide (lane 6).

Anti-SHP1 (C-terminal region) Antibody - Background

SHP1 (PTP1C, SH-PTP1, or HCP) is a protein-tyrosine phosphatase (PTP) involved in cell migration, cell proliferation, and immune cell function. This phosphatase contains two N-terminal SH2 domains and a C-terminal phosphatase domain. SHP1 associates with a variety of cytokine and growth factor receptors and regulates signal transduction through dephosphorylation of these receptors or their downstream effectors. Downstream of receptor activation, SHP1 regulates the transcriptional activity stimulated by JAK/Stat and MAPK pathways. SHP1 activity is regulated by both tyrosine and serine phosphorylation. Phosphorylation of Tyr-536 and Tyr-564 stimulates phosphatase activity and promotes interaction with Grb-2. Serine phosphorylation at Ser-591 is mediated by PKC α and leads to inhibition of phosphatase activity. Thus, phosphorylation at tyrosine relative to serine residues may be regulated by different cell signaling pathways to control SHP1 activity.