

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody

Catalog # AN1969

Specification

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody - Product Information

Primary Accession Reactivity Host Clonality Isotype Calculated MW <u>08Cl15</u> Bovine Rabbit Rabbit Polyclonal IgG 42443

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody - Additional Information

Gene ID Other Names SK1, SPHK1, SPK1 20698

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody - Images





Western blot image of recombinant his-tagged human SK1 protein. Blots were probed with rabbit polyclonals anti-SK1 (Central Region) (SP1621) at 1:1000 (lane 1) and 1:4000 (lane 2), and anti-SK1 (C-terminal region) (SP5421) at 1:1000 (lane 3) and 1:4000 (lane 4).

Anti-Sphingosine Kinase 1 (C-terminal region) Antibody - Background

Sphingolipids are metabolized into bioactive products that include ceramide, sphingosine, and sphingosine-1-phosphate (S1P). Sphingosine Kinase (SK) catalyzes the phosphorylation of the lipid sphingosine, creating S1P. S1P subsequently signals through cell surface G protein-coupled receptors, as well as intracellularly, to modulate cell proliferation, survival, motility and differentiation. Two isoforms of SK have been identified, SK1 and SK2. The mRNA for both of these isoforms is widely expressed with SK1 expression highest in brain, heart, kidney, thymus, spleen and lung, while SK2 is highest in kidney and liver. SKs can be activated through growth factor, G protein-coupled, and immunoglobulin receptor signalling. SK1 has been shown to mediate cell growth, prevention of apoptosis, and cellular transformation, and is upregulated in a variety of human tumors. Regulation of SK1 may occur through ERK mediated phosphorylation of Ser-225. This phosphorylation leads to increased activity and translocation to the plasma membrane.