

**Anti-VASP (C-terminal region) Antibody**  
**Catalog # AN2006****Specification**

---

**Anti-VASP (C-terminal region) Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P50552</a>
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Calculated MW	39830

**Anti-VASP (C-terminal region) Antibody - Additional Information**

Gene ID 7408

**Other Names**

vasodilator-stimulated phosphoprotein

**Dilution**

WB~~1:1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-VASP (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


**Shipping**


Blue Ice

**Anti-VASP (C-terminal region) Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-VASP (C-terminal region) Antibody - Images**



Western blot image of human A431 cells stimulated with calyculin A (100 nM) for 30 min. The blots were untreated (lanes 1 & 3) or treated with lambda phosphatase (lanes 2 & 4), then probed with mouse monoclonal VASP (C-term.) antibody (lanes 1 & 2) or rabbit polyclonal VASP (Thr-278) phospho-specific antibody (lanes 3 & 4).



Immunocytochemical labeling of VASP in aldehyde-fixed and NP-40-permeabilized A431 cells. The cells were labeled with mouse monoclonal VASP (C-terminal region) antibody, then the antibody was detected using appropriate secondary antibody conjugated to DyLight® 594.

### Anti-VASP (C-terminal region) Antibody - Background

Actin filament tethering and bundling are important mechanisms involved in actin superstructure assembly. The ENA/VASP family includes VASP, mena, and Ena-Vasp-like (EVL). These multidomain proteins localize to the leading edge of filopodia where they associate with AFs, interact with profilin, and compete with capping proteins at the barbed end of AFs. Artificial relocation of VASP from the plasma membrane to mitochondrial membranes inhibits filopodial formation and axon branching, while deletion of all three ENA/VASP proteins produces defects in cortical axon-tract formation. Regulation of VASP protein activity occurs through phosphorylation at Ser-157, Ser-239, and Thr-278. AMPK phosphorylates Thr-278, leading to impaired actin stress fiber assembly and changes in cell morphology.