

Anti-Vav3 Antibody

Catalog # AN2012

Specification

Anti-Vav3 Antibody - Product Information

<u>09UKW4</u>
Bovine
Rabbit
Rabbit Polyclonal
IgG
97776

Anti-Vav3 Antibody - Additional Information

Gene ID 10451 Other Names VAV3, Guanine nucleotide exchange factor VAV3

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Anti-Vav3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

Anti-Vav3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Vav3 Antibody - Images



Western blot of human Jurkat (lanes 1 & 4), HUVEC (lanes 2 & 5), and A431 (lanes 3 & 6) cells. The blots were probed with anti-Vav2 (a.a. 309-322) at a dilution of 1:500 (lanes 1-3) and anti-Vav3 (a.a. 293-305) at 1:500 (lanes 4-6).

Anti-Vav3 Antibody - Background

The Vav family of Rho-guanine nucleotide exchange factors, Vav1, Vav2, and Vav3, have central roles in transducing signals from cell surface receptors, such as integrin, growth factor and immune cell receptors to the cytoskeleton. This role includes receptor-mediated changes in the actin cytoskeleton and cell motility. Vav1 expression is normally restricted to hematopoietic cells, while Vav2 and Vav3 are more widely expressed. All three Vav isoforms have been shown to be abnormally expressed in several types of cancer. Vavs are composed of multiple domains, including a Dbl homology domain, a calponin homology domain, an acidic amino acid region, a pleckstrin homology domain, a cysteine-rich domain, and SH3 and SH2 domains. Vav activity is regulated by the phosphorylation status of several conserved tyrosine residues in the acidic region (In Vav2: Tyr-142, Tyr-159, and Tyr-172). These tyrosine residues are able to participate in autoinhibitory interactions with the Dbl homology domain and this interaction is prevented after phosphorylation of these sites leading to activation of Vav GEF activity.