

Anti-HCV Core Protein Antibody
Catalog # AN2039**Specification**

Anti-HCV Core Protein Antibody - Product InformationHost
Clonality
Isotype**Rabbit**
Rabbit Polyclonal
IgG**Anti-HCV Core Protein Antibody - Additional Information****Other Names**

hepatitis C virus

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-HCV Core Protein Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice

Anti-HCV Core Protein Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HCV Core Protein Antibody - Images**Anti-HCV Core Protein Antibody - Background**

The hepatitis C virus (HCV) core protein represents the first 191 amino acids of the viral precursor polyprotein and is cotranslationally inserted into the membrane of the endoplasmic reticulum. Hepatitis C virus (HCV) core is a viral structural protein; it also participates in some cellular processes, including transcriptional regulation. However the mechanisms of core-mediated transcriptional regulation remain poorly understood. Hepatitis C virus (HCV) core protein is thought to contribute to HCV pathogenesis through its interaction with various signal transduction pathways. In addition, HCV core antigen is a recently developed marker of hepatitis C infection. The

HCV core protein has been previously shown to circulate in the bloodstream of HCV-infected patients and inhibit host immunity through an interaction with gC1qR. Hepatitis C Virus is a positive, single stranded RNA virus in the Flaviviridae family. The genome is approximately 10,000 nucleotides and encodes a single polyprotein of about 3,000 amino acids. The polyprotein is processed by host cell and viral proteases into three major structural proteins and several non structural proteins necessary for viral replication. Hepatitis C virus (HCV) causes most cases of non-A, non-B hepatitis and results in most HCV infected people developing chronic infections, liver cirrhosis and hepatocellular carcinoma. T cell responses, including interferon-gamma production are severely suppressed in chronic HCV patients.