

Anti-Notch 2 Homolog Antibody
Catalog # AN2187**Specification**

Anti-Notch 2 Homolog Antibody - Product Information

Primary Accession	Q04721
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	265405

Anti-Notch 2 Homolog Antibody - Additional Information

Gene ID **4853**

Other Names

Neurogenic locus notch homolog protein 2, hN2

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Notch 2 Homolog Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice

Anti-Notch 2 Homolog Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Notch 2 Homolog Antibody - Images**Anti-Notch 2 Homolog Antibody - Background**

Notch signaling plays a key role in the normal development of many tissues and cell types, through diverse effects on differentiation, survival, and/or proliferation that are highly dependent on signal strength and cellular context. Members of the Notch gene family encode transmembrane receptors that are critical for various cell fate decisions. Notch family members share structural

characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Multiple human notch proteins (NOTCH1, NOTCH2, NOTCH3 and NOTCH4) have been identified and they function as receptors for membrane bound ligands. Notch signaling is also linked to tumorigenesis as first demonstrated by the identification of a recurrent t(7;9)(q34;q34.3) chromosomal translocation involving the human Notch1 gene that is found in a small subset of human pre-T-cell acute lymphoblastic leukemias (T-ALL). Since this discovery, aberrant Notch signaling has been suggested to be involved in a wide variety of human neoplasms.