

IL-8 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1013a**Specification**

IL-8 Antibody - Product Information

Application	WB, E
Primary Accession	P10145
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1.kappa

Description

Interleukin-8 (IL-8) is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. IL-8 is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. IL-8 and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q.

Immunogen

Purified recombinant fragment of human IL-8 expressed in E. Coli.

Formulation

Purified antibody in PBS containing 0.03% sodium azide.

IL-8 Antibody - Additional Information

Gene ID 3576

Other Names

Interleukin-8, IL-8, C-X-C motif chemokine 8, Chemokine (C-X-C motif) ligand 8, Emotakin, Granulocyte chemotactic protein 1, GCP-1, Monocyte-derived neutrophil chemotactic factor, MDNCF, Monocyte-derived neutrophil-activating peptide, MONAP, Neutrophil-activating protein 1, NAP-1, Protein 3-10C, T-cell chemotactic factor, MDNCF-a, GCP/IL-8 protein IV, IL8/NAP1 form I, Interleukin-8, (Ala-IL-8)77, GCP/IL-8 protein II, IL-8(1-77), IL8/NAP1 form II, MDNCF-b, IL-8(5-77), IL-8(6-77), (Ser-IL-8)72, GCP/IL-8 protein I, IL8/NAP1 form III, Lymphocyte-derived neutrophil-activating factor, LYNAF, MDNCF-c, Neutrophil-activating factor, NAF, IL-8(7-77), GCP/IL-8 protein V, IL8/NAP1 form IV, IL-8(8-77), GCP/IL-8 protein VI, IL8/NAP1 form V, IL-8(9-77), GCP/IL-8 protein III, IL8/NAP1 form VI, CXCL8, IL8

Dilution

WB~~1/500 - 1/2000

E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL-8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

IL-8 Antibody - Protein Information

Name CXCL8

Synonyms IL8

Function

Chemotactic factor that mediates inflammatory response by attracting neutrophils, basophils, and T-cells to clear pathogens and protect the host from infection (PubMed: [18692776](http://www.uniprot.org/citations/18692776), PubMed: [7636208](http://www.uniprot.org/citations/7636208)). Also plays an important role in neutrophil activation (PubMed: [2145175](http://www.uniprot.org/citations/2145175), PubMed: [9623510](http://www.uniprot.org/citations/9623510)). Released in response to an inflammatory stimulus, exerts its effect by binding to the G-protein-coupled receptors CXCR1 and CXCR2, primarily found in neutrophils, monocytes and endothelial cells (PubMed: [1840701](http://www.uniprot.org/citations/1840701), PubMed: [1891716](http://www.uniprot.org/citations/1891716)). G-protein heterotrimer (alpha, beta, gamma subunits) constitutively binds to CXCR1/CXCR2 receptor and activation by IL8 leads to beta and gamma subunits release from Galpha (GNAI2 in neutrophils) and activation of several downstream signaling pathways including PI3K and MAPK pathways (PubMed: [11971003](http://www.uniprot.org/citations/11971003), PubMed: [8662698](http://www.uniprot.org/citations/8662698)).


Cellular Location

Secreted.

IL-8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL-8 Antibody - Images




Figure 1: Western blot analysis using IL8 mouse mAb against IL8 recombinant protein.

IL-8 Antibody - References

1. Baggiolini M. et al. 1991. FEBS Lett. 307:97-101.
2. Mukaida N. et al. 1989. J.Immunol.143:1366-1371.