

PBEF1 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1157a**Specification**

PBEF1 Antibody - Product Information

Application	WB, E
Primary Accession	P43490
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

Description

PBEF1: nicotinamide phosphoribosyltransferase. This gene encodes a protein that catalyzes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, one step in the biosynthesis of nicotinamide adenine dinucleotide. The protein is an adipokine that is localized to the bloodstream and has various functions, including the promotion of vascular smooth muscle cell maturation and inhibition of neutrophil apoptosis. It also activates insulin receptor and has insulin-mimetic effects, lowering blood glucose and improving insulin sensitivity. The protein is highly expressed in visceral fat and serum levels of the protein correlate with obesity.

Immunogen

Purified recombinant fragment of PBEF1 (aa338-479) expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

PBEF1 Antibody - Additional Information

Gene ID 10135

Other Names

Nicotinamide phosphoribosyltransferase, NAMPRTase, Nampt, 2.4.2.12, Pre-B-cell colony-enhancing factor 1, Pre-B cell-enhancing factor, Visfatin, NAMPT, PBEF, PBEF1

Dilution

WB~~1/500 - 1/2000

E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PBEF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PBEF1 Antibody - Protein Information

Name NAMPT**Synonyms** PBEF, PBEF1**Function**

Catalyzes the condensation of nicotinamide with 5- phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, an intermediate in the biosynthesis of NAD. It is the rate limiting component in the mammalian NAD biosynthesis pathway. The secreted form behaves both as a cytokine with immunomodulating properties and an adipokine with anti-diabetic properties, it has no enzymatic activity, partly because of lack of activation by ATP, which has a low level in extracellular space and plasma. Plays a role in the modulation of circadian clock function. NAMPT-dependent oscillatory production of NAD regulates oscillation of clock target gene expression by releasing the core clock component: CLOCK-BMAL1 heterodimer from NAD-dependent SIRT1- mediated suppression (By similarity).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q99KQ4}. Secreted Note=Under non-inflammatory conditions, visfatin predominantly exhibits a granular pattern within the nucleus. Secreted by endothelial cells upon IL-1beta stimulation. Abundantly secreted in milk, reaching 100- fold higher concentrations compared to maternal serum


Tissue Location

Expressed in large amounts in bone marrow, liver tissue, and muscle. Also present in heart, placenta, lung, and kidney tissues

PBEF1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PBEF1 Antibody - Images




Figure 1: Western blot analysis using PBEF1 mouse mAb against truncated PBEF1-His recombinant protein (1) and full-length GFP-PBEF1(aa1-491) transfected COS7 cell lysate (2).

PBEF1 Antibody - References

1. Mol Cell Biol. 1994 Feb;14(2):1431-7. 2. Eur J Immunol. 2002 Nov;32(11):3225-34.