

RTN3 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1307a**Specification**

RTN3 Antibody - Product Information

Application	ICC, E
Primary Accession	O95197
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Calculated MW	113kDa KDa

Description

RTN3, reticulon 3. The reticulons are a group of highly conserved genes with preferential expression in neuroendocrine tissues. RTN3 may be involved in membrane trafficking in the early secretory pathway. Inhibits BACE1 activity and amyloid precursor protein processing. May induce caspase-8 cascade and apoptosis. May favor BCL2 translocation to the mitochondria upon endoplasmic reticulum stress. In case of enteroviruses infection, RTN3 may be involved in the viral replication or pathogenesis. There are 5 isoforms.

Immunogen

Purified recombinant fragment of RTN3 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

RTN3 Antibody - Additional Information

Gene ID 10313

Other Names

Reticulon-3, Homolog of ASY protein, HAP, Neuroendocrine-specific protein-like 2, NSP-like protein 2, Neuroendocrine-specific protein-like II, NSP-like protein II, NSPLII, RTN3, ASYIP, NSPL2

Dilution

ICC~~N/A

E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RTN3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RTN3 Antibody - Protein Information

Name RTN3

Synonyms ASYIP, NSPL2

Function

May be involved in membrane trafficking in the early secretory pathway. Inhibits BACE1 activity and amyloid precursor protein processing. May induce caspase-8 cascade and apoptosis. May favor BCL2 translocation to the mitochondria upon endoplasmic reticulum stress. Induces the formation of endoplasmic reticulum tubules (PubMed:25612671). Also acts as an inflammation-resolving regulator by interacting with both TRIM25 and RIGI, subsequently impairing RIGI 'Lys-63'-linked polyubiquitination leading to IRF3 and NF-kappa-B inhibition.

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein

Tissue Location

Isoform 3 is widely expressed, with highest levels in brain, where it is enriched in neuronal cell bodies from gray matter (at protein level). Three times more abundant in macula than in peripheral retina. Isoform 1 is expressed at high levels in brain and at low levels in skeletal muscle. Isoform 2 is only found in melanoma

RTN3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RTN3 Antibody - Images




Figure 1: Confocal immunofluorescence analysis of Hela (A), A431 (B) and THP-1 (C) cells using RTN3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.




Figure 1: Western blot analysis using NTRK3 mouse mAb against extracellular domain of human NTRK3 (aa32-429).

RTN3 Antibody - References

1. Biochem Biophys Res Commun. 2005 Sep 9;334(4):1198-205.
2. Brain Res Mol Brain Res. 2005 Aug 18;138(2):236-43.
3. FASEB J. 2003 Jul;17(10):1238-47.