

BDH1 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1356a

Specification

BDH1 Antibody - Product Information

Application WB, IHC, E **Primary Accession** Q02338

Reactivity Human, Mouse

Host Mouse Clonality **Monoclonal** Isotype laG1 38kDa KDa Calculated MW

Description

BDH1 (3-hydroxybutyrate dehydrogenase, type 1), it is a member of the short-chain dehydrogenase/reductase gene family. This protien forms a homotetrameric lipid-requiring enzyme of the mitochondrial membrane and has a specific requirement for phosphatidylcholine for optimal enzymatic activity. It catalyzes the interconversion of acetoacetate and (R)-3-hydroxybutyrate, the two major ketone bodies produced during fatty acid catabolism.

Immunogen

Purified recombinant fragment of human BDH1 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

BDH1 Antibody - Additional Information

Gene ID 622

Other Names

D-beta-hydroxybutyrate dehydrogenase, mitochondrial, BDH, 1.1.1.30, 3-hydroxybutyrate dehydrogenase, Short chain dehydrogenase/reductase family 9C member 1, BDH1, BDH, SDR9C1

Dilution

WB~~1/500 - 1/2000 IHC~~1/500 - 1/2000 E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BDH1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BDH1 Antibody - Protein Information



Name BDH1 (HGNC:1027)

Cellular Location

 $\label{lem:microscopic} \begin{tabular}{ll} Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q02337}. Mitochondrion matrix {ECO:0000250|UniProtKB:Q02337} \end{tabular}$

BDH1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BDH1 Antibody - Images




Figure 1: Western blot analysis using BDH1 mouse mAb against HepG2 (1) and NIH/3T3 (2) cell lysate.






Figure 2: Immunohistochemical analysis of paraffin-embedded human Prostate tissues using anti-BDH1 mouse mAb




Figure 3: Immunohistochemical analysis of paraffin-embedded human liver cancer (left) and colorectal cancer tissues (right) using BDH1 mouse mAb with DAB staining.




Figure 3: Flow cytometric analysis of Jurkat cells using CD69 mouse mAb (green) and negative control (purple).

BDH1 Antibody - References

1. Biochemistry. 1996 Jun 25;35(25):8158-65. 2. Biochemistry. 2000 Oct 3;39(39):11928-38. 3. Proc Natl Acad Sci U S A. 2001 Dec 18;98(26):15089-94.