

CDH2 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1395a

Specification

CDH2 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW **Description** WB, IHC, FC, ICC, E <u>P19022</u> Human, Mouse, Rat Mouse Monoclonal IgG1 99.8kDa KDa

This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. The protein functions during gastrulation and is required for establishment of left-right asymmetry. At certain central nervous system synapses, presynaptic to postsynaptic adhesion is mediated at least in part by this gene product.

Immunogen Purified recombinant fragment of human CDH2 expressed in E. Coli.

Formulation Ascitic fluid containing 0.03% sodium azide.

CDH2 Antibody - Additional Information

Gene ID 1000

Other Names Cadherin-2, CDw325, Neural cadherin, N-cadherin, CD325, CDH2, CDHN, NCAD

Dilution WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



CDH2 Antibody - Protein Information

Name CDH2

Synonyms CDHN, NCAD

Function

Calcium-dependent cell adhesion protein; preferentially mediates homotypic cell-cell adhesion by dimerization with a CDH2 chain from another cell. Cadherins may thus contribute to the sorting of heterogeneous cell types. Acts as a regulator of neural stem cells quiescence by mediating anchorage of neural stem cells to ependymocytes in the adult subependymal zone: upon cleavage by MMP24, CDH2-mediated anchorage is affected, leading to modulate neural stem cell quiescence. Plays a role in cell-to-cell junction formation between pancreatic beta cells and neural crest stem (NCS) cells, promoting the formation of processes by NCS cells (By similarity). Required for proper neurite branching. Required for pre- and postsynaptic organization (By similarity). CDH2 may be involved in neuronal recognition mechanism. In hippocampal neurons, may regulate dendritic spine density.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:P15116}. Cell junction. Cell surface {ECO:0000250|UniProtKB:P15116}. Cell junction, desmosome {ECO:0000250|UniProtKB:P15116}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P15116}. Note=Colocalizes with TMEM65 at the intercalated disk in cardiomyocytes. Colocalizes with OBSCN at the intercalated disk and at sarcolemma in cardiomyocytes {ECO:0000250|UniProtKB:P15116}

CDH2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CDH2 Antibody - Images






Figure 1: Western blot analysis using CDH2 mouse mAb against A431 (1), NIH/3T3 (2), Hela (3), C6 (4) and LNCap (5) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded human lung cancer (A), colon cancer (B), ovarian cancer (C) and mammary cancer(D), using CDH2 mouse mAb with DAB staining.




Figure 3: Immunofluorescence analysis of A431 cells using CDH2 mouse mAb (green). Blue:



DRAQ5 fluorescent DNA dye.




Figure 3: Immunohistochemical analysis of paraffin-embedded human Placenta tissues using CDH2 mouse mAb




Figure 4: Flow cytometric analysis of PC-2 cells using CDH2 mouse mAb (green) and negative control (purple).

CDH2 Antibody - References

1. J Biol Chem. 2007 Mar 16;282(11):8545-56. 2. Mol Cell Biochem. 2007 Aug;302(1-2):19-26. 3. Urol Oncol. 2010 Mar-Apr;28(2):180-8.