

**KLHL13 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1420a****Specification**

---

**KLHL13 Antibody - Product Information**

Application	WB, IF, IHC, FC
Primary Accession	<a href="#">Q9P2N7</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	74kDa KDa

**Description**

KLHL13 (kelch-like 13), also known as BKLHD2 (BTB and kelch domain-containing protein 2), is a 604 amino acid protein that contains six Kelch repeats and one BTB/POZ domain. Expressed predominantly in brain, KLHL13 is believed to play a role in protein ubiquitination and may function as a substrate- specific adapter of an E3 ubiquitin-protein ligase complex. E3 ligases accept a ubiquitin residue from an E2 ubiquitin-conjugating enzyme and immediately transfer that residue to a protein that is targeted for degradation. Specifically, KLHL13 interacts with KLHL9 and CUL-3, a member of the cullin family of mediators that participate in the selective targeting of proteins for ubiquitin-mediated proteolysis. Via its BTB and C-terminal Kelch (BACK) motif, KLHL13 is thought to play a role in spatially orientating substrates in the CUL-3 ligase.

**Immunogen**

Purified recombinant fragment of human KLHL13 expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**KLHL13 Antibody - Additional Information**

**Gene ID** 90293

**Other Names**

Kelch-like protein 13, BTB and kelch domain-containing protein 2, KLHL13, BKLHD2, KIAA1309

**Dilution**

WB~~1/500 - 1/2000

IF~~1/200 - 1/1000

IHC~~1/500 - 1/2000

FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

KLHL13 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## KLHL13 Antibody - Protein Information

**Name** KLHL13

**Synonyms** BKLHD2, KIAA1309

### Function

Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex required for mitotic progression and cytokinesis. The BCR(KLHL9-KLHL13) E3 ubiquitin ligase complex mediates the ubiquitination of AURKB and controls the dynamic behavior of AURKB on mitotic chromosomes and thereby coordinates faithful mitotic progression and completion of cytokinesis.

## KLHL13 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## KLHL13 Antibody - Images




Figure 1: Western blot analysis using KLHL13 mouse mAb against HeLa (1) and MCF-7 (2) cell lysate.




Figure 2: Immunofluorescence analysis of NTERA-2 cells (left) and U251 (right) cells using KLHL13 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 3: Immunohistochemical analysis of paraffin-embedded brain tissues (left) and pancreas tissues (right) using KLHL13 mouse mAb with DAB staining.




Figure 4: Flow cytometric analysis of 3T3/L1 cells using KLHL13 mouse mAb (green) and negative control (purple).

#### KLHL13 Antibody - References

1. DNA Res. 2000 Feb 28;7(1):65-73.
2. Genome Res. 2004 Sep;14(9):1711-8.
3. Cell. 2009 Jul 23;138(2):389-403.