

Rab10 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1432a

Specification

Rab10 Antibody - Product Information

WB, ICC, E Application **Primary Accession** P61026 Reactivity Human, Mouse Host Mouse Clonality Monoclonal Isotype lqG1 Calculated MW 22.5kDa KDa **Description** RAB10 belongs to the RAS (see HRAS; MIM 190020) superfamily of small GTPases. RAB proteins localize to exocytic and endocytic compartments and regulate intracellular vesicle trafficking.

Immunogen Purified recombinant fragment of human Rab10 expressed in E. Coli.

Formulation Ascitic fluid containing 0.03% sodium azide.

Rab10 Antibody - Additional Information

Gene ID 10890

Other Names Ras-related protein Rab-10, RAB10

Dilution WB~~1/500 - 1/2000 ICC~~N/A E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Rab10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Rab10 Antibody - Protein Information

Name RAB10 (<u>HGNC:9759</u>)

Function



The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes (PubMed:21248164). Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed:21248164). That Rab is mainly involved in the biosynthetic transport of proteins from the Golgi to the plasma membrane (PubMed:21248164). Regulates, for instance, SLC2A4/GLUT4 glucose transporter-enriched vesicles delivery to the plasma membrane (By similarity). In parallel, it regulates the transport of TLR4, a toll- like receptor to the plasma membrane and therefore may be important for innate immune response (By similarity). Also plays a specific role in asymmetric protein transport to the plasma membrane (PubMed:16641372). In neurons, it is involved in axonogenesis through regulation of vesicular membrane trafficking toward the axonal plasma membrane (By similarity). In epithelial cells, it regulates transport from the Golgi to the basolateral membrane (PubMed:16641372" target="_blank">16641372" target="_blank">16641372). May play a role in the basolateral recycling pathway and in phagosome maturation (By similarity). May play a role in endoplasmic reticulum dynamics and morphology controlling tubulation along microtubules and tubules fusion (PubMed:23263280). Together with LRRK2, RAB8A, and RILPL1, it regulates ciliogenesis (PubMed:30398148). When phosphorylated by LRRK2 on Thr-73, binds RILPL1 and inhibits ciliogenesis (PubMed:30398148). Participates in the export of a subset of neosynthesized proteins through a Rab8- Rab10-Rab11-dependent endososomal export route (PubMed:32344433). Targeted to and stabilized on stressed lysosomes through LRRK2 phosphorylation where it promotes the extracellular release of lysosomal content through EHBP1 and EHNP1L1 effector proteins (PubMed:30209220).

Cellular Location

Cytoplasmic vesicle membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:P24409}. Endosome membrane Recycling endosome membrane {ECO:0000250|UniProtKB:P24409}. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:P24409}. Cytoplasm, cytoskeleton, cilium basal body. Endoplasmic reticulum membrane. Cytoplasm, perinuclear region. Lysosome. Note=Associates with SLC2A4/GLUT4 storage vesicles (PubMed:22908308). Localizes to the base of the cilium when phosphorylated by LRRK2 on Thr-73 (PubMed:20576682, PubMed:30398148). Transiently associates with phagosomes (By similarity). Localizes to the endoplasmic reticulum at domains of new tubule growth (PubMed:23263280). Colocalizes with MICAL1, GRAF1/ARHGAP26 and GRAF2/ARHGAP10 on endosomal tubules (PubMed:30209220). {ECO:0000250|UniProtKB:P24409, ECO:0000269|PubMed:20576682, ECO:0000269|PubMed:22908308, ECO:0000269|PubMed:23263280,

ECO:0000269|PubMed:30209220, ECO:0000269|PubMed:30398148,

ECO:0000269 PubMed:32344433

Tissue Location

Expressed in the hippocampus (PubMed:29562525). Expressed in neutrophils (at protein level) (PubMed:29127255) Expressed in the testis (at protein level) (PubMed:28067790)

Rab10 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.



- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- <u>Cell Culture</u>

Rab10 Antibody - Images



Figure 1: Western blot analysis using Rab10 mouse mAb against Hela (1) and NIH/3T3 (2) cell lysate.



Figure 2: Immunofluorescence analysis of LOVO cells using Rab10 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.

Rab10 Antibody - References

1. Proc Natl Acad Sci U S A. 1993 Jul 15;90(14):6508-12. 2. J Biol Chem. 2003 Apr 25;278(17):15373-80. 3. Mol Med. 2010 Jul-Aug;16(7-8):247-53.