

**CHUK Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1458a****Specification**

---

**CHUK Antibody - Product Information**

Application	WB, FC, ICC, E
Primary Accession	<a href="#">O15111</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	85kDa KDa

**Description**

This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquitination pathway, thereby activating the transcription factor.

**Immunogen**

Purified recombinant fragment of human CHUK expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**CHUK Antibody - Additional Information****Gene ID 1147****Other Names**

Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, IKK-A, IKK-alpha, IKBKA, IkappaB kinase, 2.7.11.10, Conserved helix-loop-helix ubiquitous kinase, I-kappa-B kinase 1, IKK1, Nuclear factor NF-kappa-B inhibitor kinase alpha, NFKBIKA, Transcription factor 16, TCF-16, CHUK, IKKA, TCF16

**Dilution**

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

ICC~~N/A

E~~N/A

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CHUK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## CHUK Antibody - Protein Information

**Name** CHUK

**Synonyms** IKKA, TCF16

### Function

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed: <a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed: <a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed: <a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed: <a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on serine residues (PubMed: <a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed: <a href="http://www.uniprot.org/citations/35952808" target="\_blank">35952808</a>, PubMed: <a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed: <a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed: <a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed: <a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed: <a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed: <a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed: <a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed: <a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed: <a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed: <a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed: <a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). Negatively regulates the pathway by phosphorylating the scaffold protein TAXBP1 and thus promoting the assembly of the A20/TNFAIP3 ubiquitin-editing complex (composed of A20/TNFAIP3, TAX1BP1, and the E3 ligases ITCH and RNF11) (PubMed: <a href="http://www.uniprot.org/citations/21765415" target="\_blank">21765415</a>). Therefore, CHUK plays a key role in the negative feedback of NF-kappa-B canonical signaling to limit inflammatory gene activation. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes (PubMed: <a href="http://www.uniprot.org/citations/20501937" target="\_blank">20501937</a>). In turn, these complexes regulate genes encoding molecules involved in B-cell survival and lymphoid organogenesis. Also participates in the negative feedback of the non-canonical NF-kappa-B signaling pathway by phosphorylating and destabilizing MAP3K14/NIK. Within the nucleus, phosphorylates CREBBP and consequently increases both its transcriptional and histone acetyltransferase activities (PubMed: <a href="http://www.uniprot.org/citations/17434128" target="\_blank">17434128</a>). Modulates chromatin accessibility at NF-kappa-B- responsive promoters by phosphorylating histones H3 at 'Ser-10' that are subsequently acetylated at 'Lys-14' by CREBBP (PubMed: <a href="http://www.uniprot.org/citations/12789342" target="\_blank">12789342</a>). Additionally, phosphorylates the CREBBP-interacting protein NCOA3. Also phosphorylates FOXO3 and may regulate this pro-apoptotic transcription factor (PubMed: <a href="http://www.uniprot.org/citations/15084260" target="\_blank">15084260</a>). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (By similarity). Phosphorylates AMBRA1 following mitophagy induction, promoting AMBRA1 interaction with ATG8 family proteins and its mitophagic activity (PubMed: <a href="http://www.uniprot.org/citations/30217973" target="\_blank">30217973</a>).

**Cellular Location**

Cytoplasm. Nucleus Note=Shuttles between the cytoplasm and the nucleus

**Tissue Location**

Widely expressed.

**CHUK Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**CHUK Antibody - Images**

Figure 1: Western blot analysis using CHUK mouse mAb against Raji (1), Jurkat (2), THP-1 (3) and K562 (4) cell lysate.




Figure 2: Immunofluorescence analysis of NIH/3T3 cells using CHUK mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 3: Flow cytometric analysis of A549 cells using CHUK mouse mAb (blue) and negative control (red).

#### **CHUK Antibody - References**

1. Mol Cancer. 2010 Jan 5;9:1.
2. J Infect Dis. 2010 May 1;201(9):1371-80.