

**SMAD5 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1502a****Specification**

---

**SMAD5 Antibody - Product Information**

Application	WB, IHC, IF, FC
Primary Accession	<a href="#">Q99717</a>
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	52kDa KDa

**Description**

Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD). SMAD5 is required for normal development of the cardiovascular system in vivo; lack of the SMAD5 gene results in apoptosis of cardiac myocytes. 3 Upregulation of SMAD5 has been reported to mediate apoptosis of gastric epithelial cells induced by Helicobacter pylori infection. Tissue specificity: Ubiquitous.

**Immunogen**

Purified recombinant fragment of human SMAD5 expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**SMAD5 Antibody - Additional Information**

**Gene ID** 4090

**Other Names**

Mothers against decapentaplegic homolog 5, MAD homolog 5, Mothers against DPP homolog 5, JV5-1, SMAD family member 5, SMAD 5, Smad5, hSmad5, SMAD5, MADH5

**Dilution**

WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SMAD5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**SMAD5 Antibody - Protein Information**

**Name** SMAD5**Synonyms** MADH5**Function**

Transcriptional regulator that plays a role in various cellular processes including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed:<a href="http://www.uniprot.org/citations/16516194" target="\_blank">16516194</a>, PubMed:<a href="http://www.uniprot.org/citations/12064918" target="\_blank">12064918</a>). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIIs) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:<a href="http://www.uniprot.org/citations/9442019" target="\_blank">9442019</a>). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:<a href="http://www.uniprot.org/citations/33510867" target="\_blank">33510867</a>). Non-phosphorylated SMAD5 has a cytoplasmic role in energy metabolism regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed:<a href="http://www.uniprot.org/citations/28675158" target="\_blank">28675158</a>). Mechanistically, interacts with hexokinase 1/HK1 and thereby accelerates glycolysis (PubMed:<a href="http://www.uniprot.org/citations/28675158" target="\_blank">28675158</a>).

**Cellular Location**

Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4


**Tissue Location**

Ubiquitous.

**SMAD5 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SMAD5 Antibody - Images**




Figure 1: Western blot analysis using SMAD5 mouse mAb against Hela (1), SK-N-SH (2), PC-12 (3), Jurkat (4), and K562 (5) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded brain tissues (left) and lung cancer tissues (right) using SMAD5 mouse mAb with DAB staining.




Figure 3: Immunofluorescence analysis of NTERA-2 cells using SMAD5 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 4: Flow cytometric analysis of Jurkat cells using SMAD5 mouse mAb (green) and negative control (purple).

#### **SMAD5 Antibody - References**

1. Proc Natl Acad Sci U S A. 2008 Mar 11;105(10):3927-32.
2. Nat Cell Biol. 2008 May;10(5):567-74.