

JUP Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1620a**Specification****JUP Antibody - Product Information**

Application	WB, IHC, ICC, E
Primary Accession	P14923
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	82kDa KDa

Description

This gene encodes a major cytoplasmic protein which is the only known constituent common to submembranous plaques of both desmosomes and intermediate junctions. This protein forms distinct complexes with cadherins and desmosomal cadherins and is a member of the catenin family since it contains a distinct repeating amino acid motif called the armadillo repeat. Mutation in this gene has been associated with Naxos disease. Alternative splicing occurs in this gene; however, not all transcripts have been fully described.

Immunogen

Purified recombinant fragment of human JUP expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

JUP Antibody - Additional Information

Gene ID 3728

Other Names

Junction plakoglobin, Catenin gamma, Desmoplakin III, Desmoplakin-3, JUP, CTNNG, DP3

Dilution

WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
ICC~~N/A
E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

JUP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

JUP Antibody - Protein Information

Name JUP ([HGNC:6207](#))

Function

Common junctional plaque protein. The membrane-associated plaques are architectural elements in an important strategic position to influence the arrangement and function of both the cytoskeleton and the cells within the tissue. The presence of plakoglobin in both the desmosomes and in the intermediate junctions suggests that it plays a central role in the structure and function of submembranous plaques. Acts as a substrate for VE-PTP and is required by it to stimulate VE-cadherin function in endothelial cells. Can replace beta-catenin in E-cadherin/catenin adhesion complexes which are proposed to couple cadherins to the actin cytoskeleton (By similarity).

Cellular Location

Cell junction, adherens junction. Cell junction, desmosome. Cytoplasm, cytoskeleton. Cell membrane; Peripheral membrane protein. Cytoplasm {ECO:0000250|UniProtKB:Q9PVF7}. Cell junction {ECO:0000250|UniProtKB:Q9PVF7}. Nucleus {ECO:0000250|UniProtKB:Q9PVF7} Note=Cytoplasmic in a soluble and membrane-associated form. Colocalizes with DSG4 at desmosomes (PubMed:21495994)


Tissue Location

Expressed in the heart (at protein level).

JUP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using JUP mAb against human JUP (AA: 534-740) recombinant protein. (Expected MW is 48.5 kDa)




Figure 2: Western blot analysis using JUP mouse mAb against T47D (1), MCF-7 (2), SKBR-3 (3), A431 (4) and HEK293 (5) cell lysate.




Figure 3: Immunohistochemical analysis of paraffin-embedded rectum tissues using JUP mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded stomach cancer tissues using JUP mouse mAb with DAB staining.




Figure 5: Immunofluorescence analysis of U251 cells using JUP mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.

JUP Antibody - References

1. Cell. 2009 Jul 23;138(2):389-403.
2. Cancer Res. 2009 Jul 15;69(14):5734-42.