

**CCT2 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1624a****Specification****CCT2 Antibody - Product Information**

Application	E, WB, IF, FC
Primary Accession	<a href="#">P78371</a>
Reactivity	Human, Mouse, Rat, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	54kDa KDa

**Description**

The protein encoded by this gene is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Two transcript variants encoding different isoforms have been found for this gene.

**Immunogen**

Purified recombinant fragment of human CCT2 expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**CCT2 Antibody - Additional Information**

**Gene ID** 10576

**Other Names**

T-complex protein 1 subunit beta, TCP-1-beta, CCT-beta, CCT2, 99D8.1, CCTB

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CCT2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CCT2 Antibody - Protein Information**

**Name** CCT2

**Synonyms** 99D8.1, CCTB

**Function**

Component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis (PubMed:<a href="http://www.uniprot.org/citations/25467444" target="\_blank">25467444</a>). The TRiC complex mediates the folding of WRAP53/TCAB1, thereby regulating telomere maintenance (PubMed:<a href="http://www.uniprot.org/citations/25467444" target="\_blank">25467444</a>). As part of the TRiC complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PubMed:<a href="http://www.uniprot.org/citations/20080638" target="\_blank">20080638</a>). The TRiC complex plays a role in the folding of actin and tubulin (Probable).


**Cellular Location**

Cytoplasm.

**CCT2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CCT2 mAb against human CCT2 (AA: 87-290) recombinant protein. (Expected MW is 47.9 kDa)




Figure 2: Western blot analysis using CCT2 mouse mAb against Hela (1), MCF-7 (2), Jurkat (3), T47D (4), K562 (5), A431 (6), NIH/3T3 (7), PC-12 (8) and Cos7 (9) cell lysate.




Figure 3: Immunofluorescence analysis of 3T3-L1 cells using CCT2 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 4: Flow cytometric analysis of NIH/3T3 cells using CCT2 mouse mAb (blue) and negative control (red).

#### CCT2 Antibody - References

1. J Biol Chem. 2009 May 29;284(22):14939-48.
2. Mol Cell Proteomics. 2009 Jan;8(1):157-71.