

**KCNQ1 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1647a****Specification****KCNQ1 Antibody - Product Information**

Application	WB, FC, E
Primary Accession	<a href="#">P51787</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	95kDa KDa

**Description**

This gene encodes a voltage-gated potassium channel required for repolarization phase of the cardiac action potential. This protein can form heteromultimers with two other potassium channel proteins, KCNE1 and KCNE3. Mutations in this gene are associated with hereditary long QT syndrome 1 (also known as Romano-Ward syndrome), Jervell and Lange-Nielsen syndrome, and familial atrial fibrillation. This gene exhibits tissue-specific imprinting, with preferential expression from the maternal allele in some tissues, and biallelic expression in others. This gene is located in a region of chromosome 11 amongst other imprinted genes that are associated with Beckwith-Wiedemann syndrome (BWS), and itself has been shown to be disrupted by chromosomal rearrangements in patients with BWS. Alternatively spliced transcript variants have been found for this gene.

**Immunogen**

Purified recombinant fragment of human KCNQ1 expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**KCNQ1 Antibody - Additional Information**

**Gene ID** 3784

**Other Names**

Potassium voltage-gated channel subfamily KQT member 1, IKs producing slow voltage-gated potassium channel subunit alpha KvLQT1, KQT-like 1, Voltage-gated potassium channel subunit Kv7.1, KCNQ1, KCNA8, KCNA9, KVLQT1

**Dilution**

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Precautions

KCNQ1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## KCNQ1 Antibody - Protein Information

**Name** KCNQ1 ([HGNC:6294](#))

### Function

Pore-forming subunit of the voltage-gated potassium (Kv) channel involved in the regulation of cardiomyocyte excitability and important in normal development and functions of myocardium, inner ear, stomach and colon (PubMed:<a href="http://www.uniprot.org/citations/10646604" target="\_blank">10646604</a>, PubMed:<a href="http://www.uniprot.org/citations/25441029" target="\_blank">25441029</a>). Associates with KCNE beta subunits that modulates current kinetics (PubMed:<a href="http://www.uniprot.org/citations/10646604" target="\_blank">10646604</a>, PubMed:<a href="http://www.uniprot.org/citations/11101505" target="\_blank">11101505</a>, PubMed:<a href="http://www.uniprot.org/citations/19687231" target="\_blank">19687231</a>, PubMed:<a href="http://www.uniprot.org/citations/8900283" target="\_blank">8900283</a>, PubMed:<a href="http://www.uniprot.org/citations/9108097" target="\_blank">9108097</a>, PubMed:<a href="http://www.uniprot.org/citations/9312006" target="\_blank">9312006</a>). Induces a voltage-dependent current by rapidly activating and slowly deactivating potassium-selective outward current (PubMed:<a href="http://www.uniprot.org/citations/10646604" target="\_blank">10646604</a>, PubMed:<a href="http://www.uniprot.org/citations/11101505" target="\_blank">11101505</a>, PubMed:<a href="http://www.uniprot.org/citations/25441029" target="\_blank">25441029</a>, PubMed:<a href="http://www.uniprot.org/citations/8900283" target="\_blank">8900283</a>, PubMed:<a href="http://www.uniprot.org/citations/9108097" target="\_blank">9108097</a>, PubMed:<a href="http://www.uniprot.org/citations/9312006" target="\_blank">9312006</a>). Also promotes a delayed voltage activated potassium current showing outward rectification characteristic (By similarity). During beta-adrenergic receptor stimulation, participates in cardiac repolarization by associating with KCNE1 to form the I(Ks) cardiac potassium current that increases the amplitude and slows down the activation kinetics of outward potassium current I(Ks) (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/10646604" target="\_blank">10646604</a>, PubMed:<a href="http://www.uniprot.org/citations/11101505" target="\_blank">11101505</a>, PubMed:<a href="http://www.uniprot.org/citations/8900283" target="\_blank">8900283</a>, PubMed:<a href="http://www.uniprot.org/citations/9108097" target="\_blank">9108097</a>, PubMed:<a href="http://www.uniprot.org/citations/9312006" target="\_blank">9312006</a>). Muscarinic agonist oxotremorine-M strongly suppresses KCNQ1/KCNE1 current (PubMed:<a href="http://www.uniprot.org/citations/10713961" target="\_blank">10713961</a>). When associated with KCNE3, forms the potassium channel that is important for cyclic AMP-stimulated intestinal secretion of chloride ions (PubMed:<a href="http://www.uniprot.org/citations/10646604" target="\_blank">10646604</a>). This interaction with KCNE3 is reduced by 17beta-estradiol, resulting in the reduction of currents (By similarity). During conditions of increased substrate load, maintains the driving force for proximal tubular and intestinal sodium ions absorption, gastric acid secretion, and cAMP-induced jejunal chloride ions secretion (By similarity). Allows the provision of potassium ions to the luminal membrane of the secretory canaliculus in the resting state as well as during stimulated acid secretion (By similarity). When associated with KCNE2, forms a heterooligomer complex leading to currents with an apparently instantaneous activation, a rapid deactivation process and a linear current-voltage relationship and decreases the amplitude of the outward current (PubMed:<a href="http://www.uniprot.org/citations/11101505" target="\_blank">11101505</a>). When associated with KCNE4, inhibits voltage-gated potassium channel activity (PubMed:<a href="http://www.uniprot.org/citations/19687231" target="\_blank">19687231</a>). When associated with KCNE5, this complex only conducts current upon strong and continued depolarization (PubMed:<a href="http://www.uniprot.org/citations/12324418" target="\_blank">12324418</a>). Also forms a heterotetramer with KCNQ5; has a voltage-gated potassium channel activity (PubMed:<a href="http://www.uniprot.org/citations/12324418" target="\_blank">12324418</a>).

href="http://www.uniprot.org/citations/24855057" target="\_blank">24855057</a>). Binds with phosphatidylinositol 4,5-bisphosphate (PubMed:<a href="http://www.uniprot.org/citations/25037568" target="\_blank">25037568</a>). KCNQ1-KCNE2 channel associates with Na(+)-coupled myo-inositol symporter in the apical membrane of choroid plexus epithelium and regulates the myo- inositol gradient between blood and cerebrospinal fluid with an impact on neuron excitability (By similarity).

### Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane Early endosome. Membrane raft. Endoplasmic reticulum Basolateral cell membrane. Apical cell membrane {ECO:0000250|UniProtKB:P97414}; Multi-pass membrane protein. Note=Colocalized with KCNE3 at the plasma membrane (PubMed:10646604). Upon 17beta-oestradiol treatment, colocalizes with RAB5A at early endosome (PubMed:23529131). Heterotetramer with KCNQ5 is highly retained at the endoplasmic reticulum and is localized outside of lipid raft microdomains (PubMed:24855057). During the early stages of epithelial cell polarization induced by the calcium switch, it is removed from the plasma membrane to the endoplasmic reticulum, where it is retained, and redistributed to the basolateral cell surface in a PI3K-dependent manner at a later stage (PubMed:21228319). Colocalizes with SLC5A3 at the apical membrane of choroid plexus epithelium {ECO:0000250|UniProtKB:P97414, ECO:0000269|PubMed:10646604, ECO:0000269|PubMed:21228319, ECO:0000269|PubMed:23529131, ECO:0000269|PubMed:24855057}


### Tissue Location

Abundantly expressed in heart, pancreas, prostate, kidney, small intestine and peripheral blood leukocytes. Less abundant in placenta, lung, spleen, colon, thymus, testis and ovaries

### KCNQ1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using KCNQ1 mAb against human KCNQ1 (AA: 229-347) recombinant protein. (Expected MW is 74.7 kDa)




Figure 2: Western blot analysis using KCNQ1 mouse mAb against MCF-7 (1) and A431 (2) cell lysate.




Figure 3: Flow cytometric analysis of MCF-7 cells using KCNQ1 mouse mAb (green) and negative control (red).

**KCNQ1 Antibody - References**

1. Biochem Biophys Res Commun. 2009 May 29;383(2):206-9. 2. J Biol Chem. 2009 Jun 12;284(24):16452-62.