

human Splunc2 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1681a**Specification****human Splunc2 Antibody - Product Information**

Application	E, WB, FC
Primary Accession	Q96DR5
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	27kDa KDa

Description

SPLUNC2 is one member of PLUNC family, the gene undergoes alternative splicing using two 5' non-coding exons, suggesting that the gene is regulated by alternative promoters. multiple SPLUNC2 isoforms are found in the oral cavity and suggest that these proteins may be differentially regulated in distinct tissues where they may function in the innate immune response. Mucin plugs, mucous and intermediate cells of mucoepidermoid carcinomas were positive for LPLUNC1 and SPLUNC2, but areas composed of epidermoid and clear cells were negative for all PLUNCs. Papillary cystadenocarcinoma was positive for all PLUNCs.

Immunogen

Purified recombinant fragment of human Splunc2 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

human Splunc2 Antibody - Additional Information

Gene ID 140683

Other Names

BPI fold-containing family A member 2, Parotid secretory protein, PSP, Short palate, lung and nasal epithelium carcinoma-associated protein 2, BPIFA2, C20orf70, SPLUNC2

Dilution

E~~1/10000
WB~~1/500 - 1/2000
FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

human Splunc2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

human Splunc2 Antibody - Protein Information

Name BPIFA2

Synonyms C20orf70, SPLUNC2

Function

Has strong antibacterial activity against *P. aeruginosa*.

Cellular Location

Secreted.


Tissue Location

Detected in submandibular gland. Secreted into saliva.

human Splunc2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using human Splunc2 mAb against human Splunc2 (AA: 16-250) recombinant protein. (Expected MW is 50.7 kDa)




Figure 2: Flow cytometric analysis of MCF-7 cells using human Splunc2 mouse mAb (green) and negative control (red).

human Splunc2 Antibody - References

Histochem Cell Biol. 2009 Sep;132(3):339-49. Oral Dis. 2008 Oct;14(7):613-9.