

**FOS Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1710a****Specification****FOS Antibody - Product Information**

Application	WB, IHC, FC, E
Primary Accession	<a href="#">P01100</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	40.7kDa KDa

**Description**

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

**Immunogen**

Purified recombinant fragment of human FOS expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**FOS Antibody - Additional Information**

**Gene ID** 2353

**Other Names**

Proto-oncogene c-Fos, Cellular oncogene fos, G0/G1 switch regulatory protein 7, FOS, G0S7

**Dilution**

WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
FC~~1/200 - 1/400  
E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FOS Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**FOS Antibody - Protein Information**

**Name** FOS

**Synonyms** G0S7

**Function**

Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, FOS and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. On TGF-beta activation, forms a multimeric SMAD3/SMAD4/JUN/FOS complex at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation. In growing cells, activates phospholipid synthesis, possibly by activating CDS1 and PI4K2A. This activity requires Tyr-dephosphorylation and association with the endoplasmic reticulum.


**Cellular Location**

Nucleus. Endoplasmic reticulum. Cytoplasm, cytosol. Note=In quiescent cells, present in very small amounts in the cytosol. Following induction of cell growth, first localizes to the endoplasmic reticulum and only later to the nucleus. Localization at the endoplasmic reticulum requires dephosphorylation at Tyr-10 and Tyr-30

**FOS Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using FOS mAb against human FOS (AA: 116-298) recombinant protein. (Expected MW is 45.8 kDa)




Figure 2: Western blot analysis using FOS mouse mAb against HeLa (1), and HeLa (2) cell lysate.




Figure 3: Immunohistochemical analysis of paraffin-embedded colon cancer tissues using FOS mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded cerebellum tissues using FOS mouse mAb with DAB staining.




Figure 5: Flow cytometric analysis of HeLa cells using FOS mouse mAb (green) and negative control (red).

#### FOS Antibody - References

Mol Cell Biochem. 2010 Apr;337(1-2):53-63. Cancer Invest. 2009 Oct;27(8):816-24.