

**LPA Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1730a****Specification****LPA Antibody - Product Information**

Application	WB, IHC, ICC, E
Primary Accession	<a href="#">P08519.1</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	501kDa KDa

**Description**

The protein encoded by this gene is a serine proteinase that inhibits the activity of tissue-type plasminogen activator I. The encoded protein constitutes a substantial portion of lipoprotein(a) and is proteolytically cleaved, resulting in fragments that attach to atherosclerotic lesions and promote thrombogenesis. Elevated plasma levels of this protein are linked to atherosclerosis. Depending on the individual, the encoded protein contains 2-43 copies of kringle-type domains. The allele represented here contains 15 copies of the kringle-type repeats and corresponds to that found in the reference genome sequence.

**Immunogen**

Purified recombinant fragment of human LPA (AA: 4330-4521) expressed in E. Coli. <br /> <br />

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**LPA Antibody - Additional Information****Dilution**

WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
ICC~~N/A  
E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

LPA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**LPA Antibody - Protein Information**

## LPA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Figure 1: Western blot analysis using LPA mAb against human LPA recombinant protein. (Expected MW is 34.1 kDa)




Figure 2: Immunohistochemical analysis of paraffin-embedded liver cancer tissues using LPA mouse mAb with DAB staining.




Figure 3: Immunohistochemical analysis of paraffin-embedded breast cancer tissues using LPA mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using LPA mouse mAb with DAB staining.




Figure 5: Immunofluorescence analysis of HepG2 cells using LPA mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

#### LPA Antibody - References

1. J Lipid Res. 2010 Oct;51(10):3055-61.
2. Thromb Res. 2010 Sep;126(3):222-6.