

SPIB Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1754a**Specification****SPIB Antibody - Product Information**

Application	WB, IHC, FC, ICC, E
Primary Accession	Q01892
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	28.8kDa KDa

Description

The protein encoded by this gene is a transcriptional activator that binds to the PU-box (5'-GAGGAA-3') and acts as a lymphoid-specific enhancer. Four transcript variants encoding different isoforms have been found for this gene.

Immunogen

Purified recombinant fragment of human SPIB (AA: 200-252) expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

SPIB Antibody - Additional Information

Gene ID 6689

Other Names

Transcription factor Spi-B, SPIB

Dilution

WB~~1/500 - 1/2000

IHC~~1/200 - 1/1000

FC~~1/200 - 1/400

ICC~~N/A

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SPIB Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SPIB Antibody - Protein Information

Name SPIB**Function**

Sequence specific transcriptional activator which binds to the PU-box, a purine-rich DNA sequence (5'-GAGGAA-3') that can act as a lymphoid-specific enhancer. Promotes development of plasmacytoid dendritic cells (pDCs), also known as type 2 DC precursors (pre-DC2) or natural interferon (IFN)-producing cells. These cells have the capacity to produce large amounts of interferon and block viral replication. May be required for B-cell receptor (BCR) signaling, which is necessary for normal B-cell development and antigenic stimulation.

Cellular Location

[Isoform 1]: Nucleus


Tissue Location

Expressed in plasmacytoid dendritic cells (pDCs) and B-cells, not expressed in T-cells or granulocytes. May also be enriched in stem cell populations of the liver

SPIB Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using SPIB mAb against human SPIB recombinant protein. (Expected MW is 32 kDa)




Figure 2: Western blot analysis using SPIB mouse mAb against A549 (1), PC-3 (2), and NIH3T3 (3) cell lysate.




Figure 3: Immunofluorescence analysis of HeLa cells using SPIB mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 4: Flow cytometric analysis of NIH3T3 cells using SPIB mouse mAb (green) and negative control (red).




Figure 5: Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using SPIB mouse mAb with DAB staining.




Figure 6: Immunohistochemical analysis of paraffin-embedded esophageal cancer with DAB staining.

SPIB Antibody - References

1. J Gen Virol. 2010 Dec;91(Pt 12):3042-52. 2. Eur J Immunol. 2008 Sep;38(9):2389-400.