

**CSNK2A2 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1755a****Specification****CSNK2A2 Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | E, WB, FC              |
| Primary Accession | <a href="#">P19784</a> |
| Reactivity        | Human, Mouse           |
| Host              | Mouse                  |
| Clonality         | Monoclonal             |
| Isotype           | IgG1                   |
| Calculated MW     | 41.2kDa KDa            |

**Description**

Casein kinase II (CK2) is a constitutively active, ubiquitously expressed serine/threonine protein kinase that is thought to have a regulatory function in cell proliferation, cell differentiation and apoptosis. CK2 functions as a tetrameric complex consisting of two regulatory beta-subunits and two catalytic units (alpha and alpha') in a homomeric or heteromeric conformation. Whilst the alpha- and alpha'-subunits are catalytically identical, proteins that regulate CK2, such as cdc2 and Hsp90, preferentially bind to the alpha and not the alpha'-subunit. CK2 can phosphorylate a number of key intracellular signaling proteins implicated in tumor suppression (p53 and PTEN) and tumorigenesis (myc, jun, NF-kappaB). CK2 is also thought to influence Wnt signaling via beta-catenin phosphorylation and the PI 3-K signaling pathway via the phosphorylation of Akt.

**Immunogen**

Purified recombinant fragment of human CSNK2A2 (AA: 194-350) expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**CSNK2A2 Antibody - Additional Information****Gene ID** 1459**Other Names**

Casein kinase II subunit alpha', CK II alpha', 2.7.11.1, CSNK2A2, CK2A2

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CSNK2A2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## CSNK2A2 Antibody - Protein Information

**Name** CSNK2A2

**Synonyms** CK2A2

### Function

Catalytic subunit of a constitutively active serine/threonine-protein kinase complex that phosphorylates a large number of substrates containing acidic residues C-terminal to the phosphorylated serine or threonine (PubMed:<a href="http://www.uniprot.org/citations/11239457" target="\_blank">11239457</a>, PubMed:<a href="http://www.uniprot.org/citations/11704824" target="\_blank">11704824</a>, PubMed:<a href="http://www.uniprot.org/citations/16193064" target="\_blank">16193064</a>, PubMed:<a href="http://www.uniprot.org/citations/30898438" target="\_blank">30898438</a>). Regulates numerous cellular processes, such as cell cycle progression, apoptosis and transcription, as well as viral infection (PubMed:<a href="http://www.uniprot.org/citations/11704824" target="\_blank">11704824</a>, PubMed:<a href="http://www.uniprot.org/citations/16193064" target="\_blank">16193064</a>, PubMed:<a href="http://www.uniprot.org/citations/30898438" target="\_blank">30898438</a>). May act as a regulatory node which integrates and coordinates numerous signals leading to an appropriate cellular response (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). During mitosis, functions as a component of the p53/TP53-dependent spindle assembly checkpoint (SAC) that maintains cyclin-B-CDK1 activity and G2 arrest in response to spindle damage (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). Also required for p53/TP53-mediated apoptosis, phosphorylating 'Ser-392' of p53/TP53 following UV irradiation (PubMed:<a href="http://www.uniprot.org/citations/11239457" target="\_blank">11239457</a>). Phosphorylates a number of DNA repair proteins in response to DNA damage, such as MDC1, RAD9A, RAD51 and HTATSF1, promoting their recruitment to DNA damage sites (PubMed:<a href="http://www.uniprot.org/citations/20545769" target="\_blank">20545769</a>, PubMed:<a href="http://www.uniprot.org/citations/21482717" target="\_blank">21482717</a>, PubMed:<a href="http://www.uniprot.org/citations/22325354" target="\_blank">22325354</a>, PubMed:<a href="http://www.uniprot.org/citations/26811421" target="\_blank">26811421</a>, PubMed:<a href="http://www.uniprot.org/citations/30898438" target="\_blank">30898438</a>, PubMed:<a href="http://www.uniprot.org/citations/35597237" target="\_blank">35597237</a>). Can also negatively regulate apoptosis (PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). Phosphorylates the caspases CASP9 and CASP2 and the apoptotic regulator NOL3 (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). Phosphorylation protects CASP9 from cleavage and activation by CASP8, and inhibits the dimerization of CASP2 and activation of CASP8 (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). Regulates transcription by direct phosphorylation of RNA polymerases I, II, III and IV (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">19387551</a>). Also phosphorylates and regulates numerous transcription factors including NF-kappa-B, STAT1, CREB1, IRF1, IRF2, ATF1, SRF, MAX, JUN, FOS, MYC and MYB (PubMed:<a

href="http://www.uniprot.org/citations/12631575" target="\_blank">>12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">>19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">>19387551</a>). Phosphorylates Hsp90 and its co-chaperones FKBP4 and CDC37, which is essential for chaperone function (PubMed:<a href="http://www.uniprot.org/citations/19387550" target="\_blank">>19387550</a>). Regulates Wnt signaling by phosphorylating CTNNB1 and the transcription factor LEF1 (PubMed:<a href="http://www.uniprot.org/citations/19387549" target="\_blank">>19387549</a>). Acts as an ectokinase that phosphorylates several extracellular proteins (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">>12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">>19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">>19387551</a>). During viral infection, phosphorylates various proteins involved in the viral life cycles of EBV, HSV, HBV, HCV, HIV, CMV and HPV (PubMed:<a href="http://www.uniprot.org/citations/12631575" target="\_blank">>12631575</a>, PubMed:<a href="http://www.uniprot.org/citations/19387552" target="\_blank">>19387552</a>, PubMed:<a href="http://www.uniprot.org/citations/19387551" target="\_blank">>19387551</a>).

### Cellular Location

Nucleus {ECO:0000250|UniProtKB:O54833}. Cytoplasm {ECO:0000250|UniProtKB:O54833}.


Note=Interaction with SIRT6 prevents translocation into the nucleus.

{ECO:0000250|UniProtKB:O54833}

### CSNK2A2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CSNK2A2 mAb against human CSNK2A2 recombinant protein.  
(Expected MW is 44 kDa)




Figure 2: Western blot analysis using CSNK2A2 mouse mAb against HeLa (1), MCF-7 (2), HepG2 (3), Jurkat (4), NIH3T3 (5), and PC-12 (6) cell lysate.




Figure 3: Flow cytometric analysis of HepG2 cells using CSNK2A2 mouse mAb (green) and negative control (red).

#### CSNK2A2 Antibody - References

1. Eur J Immunol. 2009 Jan;39(1):267-79. 2. Cell Mol Life Sci. 2009 Jan;66(2):339-49.

