

EIF2A Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1898a**Specification****EIF2A Antibody - Product Information**

| | |
|-------------------|---------------------------|
| Application | WB, IHC, FC, E |
| Primary Accession | Q9BY44 |
| Reactivity | Human, Mouse, Rat, Monkey |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Calculated MW | 65kDa KDa |

Description

EIF2A is a 65-kD protein that catalyzes the formation of puromycin-sensitive 80S preinitiation complexes.

Immunogen

Purified recombinant fragment of human EIF2A (AA: 448-576) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide.

EIF2A Antibody - Additional Information

Gene ID 83939

Other Names

Eukaryotic translation initiation factor 2A, eIF-2A, 65 kDa eukaryotic translation initiation factor 2A, Eukaryotic translation initiation factor 2A, N-terminally processed, EIF2A

Dilution

WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
FC~~1/200 - 1/400
E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EIF2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

EIF2A Antibody - Protein Information

Name EIF2A

Function

Functions in the early steps of protein synthesis of a small number of specific mRNAs. Acts by directing the binding of methionyl- tRNAi to 40S ribosomal subunits. In contrast to the eIF-2 complex, it binds methionyl-tRNAi to 40S subunits in a codon-dependent manner, whereas the eIF-2 complex binds methionyl-tRNAi to 40S subunits in a GTP-dependent manner.

Tissue Location

Widely expressed. Expressed at higher level in pancreas, heart, brain and placenta.

EIF2A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using EIF2A mAb against human EIF2A (AA: 448-576) recombinant

protein. (Expected MW is 40.3 kDa)




Figure 2: Western blot analysis using EIF2A mAb against HEK293 (1) and EIF2A (AA: 448-576)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using EIF2A mouse mAb against MCF-7 (1), PC-12 (2), HepG2 (3), Hela (4), Cos7 (5), K562 (6), Jurkat (7), A431 (8) and NIH/3T3 (9) cell lysate.




Figure 4: Flow cytometric analysis of HepG2 cells using EIF2A mouse mAb (green) and negative control (red).




Figure 5: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using EIF2A mouse mAb with DAB staining.




Figure 6: Immunohistochemical analysis of paraffin-embedded bladder cancer tissues using EIF2A mouse mAb with DAB staining.

EIF2A Antibody - Background

The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit is one of the gamma regulatory subunits of AMPK. Alternatively spliced transcript variants encoding distinct isoforms have been observed. ;

EIF2A Antibody - References

1. Mol Biol (Mosk). 2010 Sep-Oct;44(5):859-66.
2. Cancer Res. 2009 Feb 15;69(4):1545-52.