

EIF2A Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1904a**Specification****EIF2A Antibody - Product Information**

Application	WB, IHC, FC, ICC, E
Primary Accession	Q9BY44
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	65kDa KDa

Description

EIF2A is a 65-kD protein that catalyzes the formation of puromycin-sensitive 80S preinitiation complexes

Immunogen

Purified recombinant fragment of human EIF2A (AA: 448-576) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide.

EIF2A Antibody - Additional Information

Gene ID 83939

Other Names

Eukaryotic translation initiation factor 2A, eIF-2A, 65 kDa eukaryotic translation initiation factor 2A, Eukaryotic translation initiation factor 2A, N-terminally processed, EIF2A

Dilution

WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
FC~~1/200 - 1/400
ICC~~N/A
E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EIF2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

EIF2A Antibody - Protein Information

Name EIF2A**Function**

Functions in the early steps of protein synthesis of a small number of specific mRNAs. Acts by directing the binding of methionyl- tRNAi to 40S ribosomal subunits. In contrast to the eIF-2 complex, it binds methionyl-tRNAi to 40S subunits in a codon-dependent manner, whereas the eIF-2 complex binds methionyl-tRNAi to 40S subunits in a GTP-dependent manner.


Tissue Location

Widely expressed. Expressed at higher level in pancreas, heart, brain and placenta.

EIF2A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using EIF2A mAb against human EIF2A (AA: 448-576) recombinant protein. (Expected MW is 40.3 kDa)




Figure 2: Western blot analysis using EIF2A mAb against HEK293 (1) and EIF2A (AA: 448-576)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using EIF2A mouse mAb against MCF-7 (1), PC-12 (2), HepG2 (3), Hela (4), Cos7 (5), K562 (6), Jurkat (7), A431 (8) and NIH/3T3 (9) cell lysate.




Figure 4: Immunofluorescence analysis of HepG2 cells using EIF2A mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Secondary antibody from Fisher (Cat#: 35503)




Figure 5: Flow cytometric analysis of HepG2 cells using EIF2A mouse mAb (green) and negative control (red).




Figure 6: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using EIF2A mouse mAb with DAB staining.

EIF2A Antibody - Background

This gene encodes a telomere specific protein, TERF2, which is a component of the telomere nucleoprotein complex. This protein is present at telomeres in metaphase of the cell cycle, is a second negative regulator of telomere length and plays a key role in the protective activity of

telomeres. While having similar telomere binding activity and domain organization, TERF2 differs from TERF1 in that its N terminus is basic rather than acidic. ; ;

EIF2A Antibody - References

1. Mol Biol (Mosk). 2010 Sep-Oct;44(5):859-66.
2. Cancer Res. 2009 Feb 15;69(4):1545-52.