

TIE1 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1918a**Specification****TIE1 Antibody - Product Information**

Application	WB, IHC, E
Primary Accession	P35590
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	125kDa KDa

Description

This gene encodes a member of the tyrosine protein kinase family. The encoded protein plays a critical role in angiogenesis and blood vessel stability by inhibiting angiopoietin 1 signaling through the endothelial receptor tyrosine kinase Tie2. Ectodomain cleavage of the encoded protein relieves inhibition of Tie2 and is mediated by multiple factors including vascular endothelial growth factor. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Immunogen

Purified recombinant fragment of human TIE1 (AA: 385-607) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide.

TIE1 Antibody - Additional Information

Gene ID 7075

Other Names

Tyrosine-protein kinase receptor Tie-1, 2.7.10.1, TIE1, TIE

Dilution

WB~~1/500 - 1/2000

IHC~~1/200 - 1/1000

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TIE1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TIE1 Antibody - Protein Information

Name TIE1

Synonyms TIE

Function

Transmembrane tyrosine-protein kinase that may modulate TEK/TIE2 activity and contribute to the regulation of angiogenesis.

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Specifically expressed in developing vascular endothelial cells.

TIE1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using TIE1 mAb against human TIE1 (AA: 385-607) recombinant protein. (Expected MW is 50.6 kDa)




Figure 2: Western blot analysis using TIE1 mAb against HEK293 (1) and TIE1 (AA: 385-607)-hlgGfC transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using TIE1 mouse mAb against HepG2 cell lysate.




Figure 4: Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using TIE1 mouse mAb with DAB staining.




Figure 5: Immunohistochemical analysis of paraffin-embedded kidney tissues using TIE1 mouse mAb with DAB staining.

TIE1 Antibody - Background

This locus encodes the transforming growth factor (TGF)-beta type III receptor. The encoded receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFBR3, which may inhibit TGFB signaling. Decreased expression of this receptor has been observed in various cancers. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. ; ; ;

TIE1 Antibody - References

1. Int J Oncol. 2007 Oct;31(4):893-7.
2. Cancer. 2002 Mar 1;94(5):1517-21.