

DAPK3 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1931a

Specification

DAPK3 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW **Description** WB, IHC, FC, E <u>043293</u> Human Mouse Monoclonal IgG1 52.5kDa KDa

Death-associated protein kinase 3 (DAPK3) induces morphological changes in apoptosis when overexpressed in mammalian cells. These results suggest that DAPK3 may play a role in the induction of apoptosis.

Immunogen Purified recombinant fragment of human DAPK3 (AA: 28-161) expressed in E. Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide.

DAPK3 Antibody - Additional Information

Gene ID 1613

Other Names Death-associated protein kinase 3, DAP kinase 3, 2.7.11.1, DAP-like kinase, Dlk, MYPT1 kinase, Zipper-interacting protein kinase, ZIP-kinase, DAPK3, ZIPK

Dilution WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DAPK3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DAPK3 Antibody - Protein Information



Name DAPK3

Synonyms ZIPK

Function

Serine/threonine kinase which is involved in the regulation of apoptosis, autophagy, transcription, translation and actin cytoskeleton reorganization. Involved in the regulation of smooth muscle contraction. Regulates both type I (caspase-dependent) apoptotic and type II (caspase-independent) autophagic cell deaths signal, depending on the cellular setting. Involved in regulation of starvation-induced autophagy. Regulates myosin phosphorylation in both smooth muscle and non-muscle cells. In smooth muscle, regulates myosin either directly by phosphorylating MYL12B and MYL9 or through inhibition of smooth muscle myosin phosphatase (SMPP1M) via phosphorylation of PPP1R12A; the inhibition of SMPP1M functions to enhance muscle responsiveness to Ca(2+) and promote a contractile state. Phosphorylates MYL12B in non-muscle cells leading to reorganization of actin cytoskeleton. Isoform 2 can phosphorylate myosin, PPP1R12A and MYL12B. Overexpression leads to condensation of actin stress fibers into thick bundles. Involved in actin filament focal adhesion dynamics. The function in both reorganization of actin cytoskeleton and focal adhesion dissolution is modulated by RhoD. Positively regulates canonical Wnt/beta-catenin signaling through interaction with NLK and TCF7L2. Phosphorylates RPL13A on 'Ser-77' upon interferon-gamma activation which is causing RPL13A release from the ribosome, RPL13A association with the GAIT complex and its subsequent involvement in transcript-selective translation inhibition. Enhances transcription from AR-responsive promoters in a hormone- and kinase- dependent manner. Involved in regulation of cell cycle progression and cell proliferation. May be a tumor suppressor.

Cellular Location

Nucleus. Nucleus, PML body {ECO:0000250|UniProtKB:O54784}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:O54784}. Chromosome, centromere. Cytoplasm. Cytoplasm, cytoskeleton, spindle. Midbody Note=Predominantly localizes to the cytoplasm but can shuttle between the nucleus and cytoplasm; cytoplasmic localization is promoted by phosphorylation at Thr-299 and involves Rho/Rock signaling [Isoform 2]: Nucleus. Cytoplasm

Tissue Location

Widely expressed. Isoform 1 and isoform 2 are expressed in the bladder smooth muscle.

DAPK3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>





Figure 1: Western blot analysis using DAPK3 mAb against human DAPK3 (AA: 28-161) recombinant protein. (Expected MW is 41.6 kDa)



Figure 2: Western blot analysis using DAPK3 mAb against HEK293 (1) and DAPK3 (AA: 28-161)-hlgGFc transfected HEK293 (2) cell lysate.





Figure 3: Western blot analysis using DAPK3 mouse mAb against A431 (1), K562 (2) cell lysate.



Figure 4: Flow cytometric analysis of A431 cells using DAPK3 mouse mAb (green) and negative control (red).



Figure 6: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using DAPK3 mouse mAb with DAB staining.

DAPK3 Antibody - Background

The protein encoded by this gene is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is capable of activating very long-chain fatty-acids containing 24- and 26-carbons. It is expressed in liver and associated with endoplasmic reticulum but not with peroxisomes. Its primary role is in fatty acid elongation or complex lipid synthesis rather than in degradation. This gene has a mouse



ortholog.;;

DAPK3 Antibody - References

1. Cancer Res. 2011 Apr 15;71(8):3152-61.2. Int J Cancer. 2009 Apr 1;124(7):1587-93.