

**TCL1A Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1985a****Specification****TCL1A Antibody - Product Information**

Application	WB, FC, E
Primary Accession	<a href="#">P56279</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	13.5kDa KDa

**Description**

Overexpression of the TCL1 gene in humans has been implicated in the development of mature T cell leukemia, in which chromosomal rearrangements bring the TCL1 gene in close proximity to the T-cell antigen receptor (TCR)-alpha (MIM 186880) or TCR-beta (MIM 186930) regulatory elements (summarized by Virgilio et al., 1998 [PubMed 9520462]). In normal T cells TCL1 is expressed in CD4-/CD8- cells, but not in cells at later stages of differentiation. TCL1 functions as a coactivator of the cell survival kinase AKT (MIM 164730) (Laine et al., 2000 [PubMed 10983986]).

**Immunogen**

Purified recombinant fragment of human TCL1A (AA: 10-104) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide.

**TCL1A Antibody - Additional Information**

**Gene ID** 8115

**Other Names**

T-cell leukemia/lymphoma protein 1A, Oncogene TCL-1, Oncogene TCL1, Protein p14 TCL1, TCL1A, TCL1

**Dilution**

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TCL1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**TCL1A Antibody - Protein Information**

**Name** TCL1A

**Synonyms** TCL1

**Function**

Enhances the phosphorylation and activation of AKT1, AKT2 and AKT3. Promotes nuclear translocation of AKT1. Enhances cell proliferation, stabilizes mitochondrial membrane potential and promotes cell survival.

**Cellular Location**

Cytoplasm. Nucleus. Microsome. Endoplasmic reticulum. Note=Microsomal fraction


**Tissue Location**

Restricted in the T-cell lineage to immature thymocytes and activated peripheral lymphocytes. Preferentially expressed early in T- and B-lymphocyte differentiation

**TCL1A Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using TCL1A mAb against human TCL1A (AA: 10-104) recombinant protein. (Expected MW is 37.3 kDa)




Figure 2: Western blot analysis using TCL1A mAb against HEK293 (1) and TCL1A (AA: 10-104)-hlgGfC transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using TCL1A mouse mAb (green) and negative control (red).

### **TCL1A Antibody - Background**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a protein identified as belonging to both the 28S and the 39S subunits. Alternative splicing results in multiple transcript variants. Pseudogenes corresponding to this gene are found on chromosomes 4q, 6p, 6q, 7p, and 15q. ;

### **TCL1A Antibody - References**

1. Blood. 2012 Aug 23;120(8):1613-23.2. Histopathology. 2010 Jul;57(1):152-7.