

**HDAC9 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2081a****Specification****HDAC9 Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, FC, E              |
| Primary Accession | <a href="#">O9UKV0</a> |
| Reactivity        | Human                  |
| Host              | Mouse                  |
| Clonality         | Monoclonal             |
| Isotype           | IgG1                   |
| Calculated MW     | 111.3kDa KDa           |

**Description**

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene has sequence homology to members of the histone deacetylase family. This gene is orthologous to the Xenopus and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined.

**Immunogen**

Purified recombinant fragment of human HDAC9 (AA: 343-569) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**HDAC9 Antibody - Additional Information**

**Gene ID** 9734

**Other Names**

Histone deacetylase 9, HD9, 3.5.1.98, Histone deacetylase 7B, HD7, HD7b, Histone deacetylase-related protein, MEF2-interacting transcription repressor MITR, HDAC9, HDAC7, HDAC7B, HDRP, KIAA0744, MITR

**Dilution**

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

HDAC9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **HDAC9 Antibody - Protein Information**

**Name** HDAC9

**Synonyms** HDAC7, HDAC7B, HDRP, KIAA0744, MITR

### **Function**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription.

### **Cellular Location**

Nucleus.

### **Tissue Location**

Broadly expressed, with highest levels in brain, heart, muscle and testis. Isoform 3 is present in human bladder carcinoma cells (at protein level).

## **HDAC9 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)