

# **AOF1** Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2105a

# **Specification**

## **AOF1 Antibody - Product Information**

Application WB, IHC, FC, ICC, E

Primary Accession
Reactivity
Host
Clonality
Host
Monoclonal
Isotype

ONUMB 18

Muse
Monoclonal
IgG1

Calculated MW 92kDa KDa

**Description** 

Flavin-dependent histone demethylases, such as KDM1B, regulate histone lysine methylation, an epigenetic mark that regulates gene expression and chromatin function.

### **Immunogen**

Purified recombinant fragment of human AOF1 (AA: 6-129) expressed in E. Coli.

### **Formulation**

Purified antibody in PBS with 0.05% sodium azide

# **AOF1 Antibody - Additional Information**

## Gene ID 221656

## **Other Names**

Lysine-specific histone demethylase 1B, 1.-.-., Flavin-containing amine oxidase domain-containing protein 1, Lysine-specific histone demethylase 2, KDM1B, AOF1, C6orf193, LSD2

# Dilution

WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000

# Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

AOF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **AOF1 Antibody - Protein Information**



# Name KDM1B (HGNC:21577)

### **Function**

Histone demethylase that demethylates 'Lys-4' of histone H3, a specific tag for epigenetic transcriptional activation, thereby acting as a corepressor. Required for de novo DNA methylation of a subset of imprinted genes during oogenesis. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed. Demethylates both mono- and di-methylated 'Lys-4' of histone H3. Has no effect on tri-methylated 'Lys-4', mono-, di- or tri-methylated 'Lys-9', mono-, di- or tri-methylated 'Lys-27', mono-, di- or tri-methylated 'Lys-36' of histone H3, or on mono-, di- or tri-methylated 'Lys-20' of histone H4. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of GLYR1 to achieve such activity, they form a multifunctional enzyme complex that modifies transcribed chromatin and facilitates Pol II transcription through nucleosomes (PubMed:<a href="http://www.uniprot.org/citations/30970244" target="blank">30970244</a>).

### **Cellular Location**

Nucleus. Chromosome. Note=Found in actively RNAPollI- transcribed gene bodies.

# **AOF1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture