

## **WNT3A Antibody**

Purified Mouse Monoclonal Antibody Catalog # AO2160a

## **Specification**

# **WNT3A Antibody - Product Information**

Application WB, IHC, FC, ICC, E

Primary Accession
Reactivity
Human
Host
Clonality
Honoclonal
Isotype
Reactivity
Human
Mouse
Monoclonal

Calculated MW 39.4kDa KDa

**Description** 

The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 96% amino acid identity to mouse Wnt3A protein, and 84% to human WNT3 protein, another WNT gene product. This gene is clustered with WNT14 gene, another family member, in chromosome 1q42 region.

### **Immunogen**

Purified recombinant fragment of human WNT3A (AA: 170-352) expressed in E. Coli.

#### Formulation

Purified antibody in PBS with 0.05% sodium azide

### **WNT3A Antibody - Additional Information**

Gene ID 89780

### **Other Names**

Protein Wnt-3a, WNT3A

### **Dilution**

WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

WNT3A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



# **WNT3A Antibody - Protein Information**

#### Name WNT3A

### **Function**

Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Functions in the canonical Wnt signaling pathway that results in activation of transcription factors of the TCF/LEF family (PubMed:<a href="http://www.uniprot.org/citations/20093360" target="\_blank">20093360</a>, PubMed:<a href="http://www.uniprot.org/citations/21244856" target="\_blank">21244856" target="\_blank">21244856</a>, PubMed:<a href="http://www.uniprot.org/citations/24841207" target="\_blank">24841207</a>, PubMed:<a href="http://www.uniprot.org/citations/26902720" target="\_blank">26902720</a>). Required for normal embryonic mesoderm development and formation of caudal somites. Required for normal morphogenesis of the developing neural tube (By similarity). Mediates self-renewal of the stem cells at the bottom on intestinal crypts (in vitro) (PubMed:<a href="http://www.uniprot.org/citations/26902720" target="\_blank">26902720</a>).

#### **Cellular Location**

Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P27467}. Secreted

### **Tissue Location**

Moderately expressed in placenta and at low levels in adult lung, spleen, and prostate.

### **WNT3A Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture