

FOXM1 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2180a**Specification****FOXM1 Antibody - Product Information**

Application	WB, FC, E
Primary Accession	Q08050
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	84.3kDa KDa

Description

The protein encoded by this gene is a transcriptional activator involved in cell proliferation. The encoded protein is phosphorylated in M phase and regulates the expression of several cell cycle genes, such as cyclin B1 and cyclin D1. Several transcript variants encoding different isoforms have been found for this gene.

Immunogen

Purified recombinant fragment of human FOXM1 (AA: 649-748) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

FOXM1 Antibody - Additional Information

Gene ID 2305

Other Names

Forkhead box protein M1, Forkhead-related protein FKHL16, Hepatocyte nuclear factor 3 forkhead homolog 11, HFH-11, HNF-3/fork-head homolog 11, M-phase phosphoprotein 2, MPM-2 reactive phosphoprotein 2, Transcription factor Trident, Winged-helix factor from INS-1 cells, FOXM1, FKHL16, HFH11, MPP2, WIN

Dilution

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FOXM1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FOXM1 Antibody - Protein Information

Name FOXM1**Synonyms** FKHL16, HFH11, MPP2, WIN**Function**

Transcription factor regulating the expression of cell cycle genes essential for DNA replication and mitosis (PubMed:19160488, PubMed:20360045). Plays a role in the control of cell proliferation (PubMed:19160488). Also plays a role in DNA break repair, participating in the DNA damage checkpoint response (PubMed:17101782). Promotes transcription of PHB2 (PubMed:33754036).

Cellular Location

Nucleus.

Tissue Location

Expressed in thymus, testis, small intestine, colon followed by ovary. Appears to be expressed only in adult organs containing proliferating/cycling cells or in response to growth factors. Also expressed in epithelial cell lines derived from tumors Not expressed in resting cells. Isoform 2 is highly expressed in testis

FOXM1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)