

Rab5a Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2187a

Specification

Rab5a Antibody - Product Information

Application E, WB, IF, FC, IHC

Primary Accession
Reactivity
Host
Clonality
Isotype
Reactivity
Human
Mouse
Monoclonal
IgG1

Calculated MW 23.7kDa KDa

Description

RAB5A (RAB5A, Member RAS Oncogene Family) is a Protein Coding gene. Diseases associated with RAB5A include borna disease and choroideremia. Among its related pathways are Ras signaling pathway and Endocytosis. GO annotations related to this gene include GTP binding and GDP binding. An important paralog of this gene is RAB5C.

Immunogen

Purified recombinant fragment of human Rab5a (AA: 1-215) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

Rab5a Antibody - Additional Information

Gene ID 5868

Other Names

Ras-related protein Rab-5A, RAB5A, RAB5

Dilution

E~~1/10000 WB~~1/500 - 1/2000 IF~~1/200 - 1/1000 FC~~1/200 - 1/400 IHC~~1/200 - 1/1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Rab5a Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Rab5a Antibody - Protein Information



Name RAB5A

Synonyms RAB5

Function

Small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate cellular responses such as of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Active GTP-bound form is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes (PubMed:<a

 $href="http://www.uniprot.org/citations/10818110" target="_blank">10818110, PubMed:14617813, PubMed:16410077, PubMed:15378032). Contributes to the regulation of filopodia extension (PubMed:15378032).$

href="http://www.uniprot.org/citations/14978216" target="_blank">14978216). Required for the exosomal release of SDCBP, CD63, PDCD6IP and syndecan (PubMed:22660413). Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 (By similarity).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane; Lipid- anchor. Melanosome. Cytoplasmic vesicle. Cell projection, ruffle {ECO:0000250|UniProtKB:P18066}. Membrane Cytoplasm, cytosol. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9CQD1}. Endosome membrane Note=Enriched in stage I melanosomes (PubMed:17081065). Alternates between membrane-bound and cytosolic forms (Probable) {ECO:0000269|PubMed:17081065, ECO:0000305}

Rab5a Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture