

#### **PDE1B Antibody**

Purified Mouse Monoclonal Antibody Catalog # AO2283a

## **Specification**

#### **PDE1B Antibody - Product Information**

Application WB, FC, E
Primary Accession Q01064

Reactivity Human, Mouse, Monkey

Host Mouse
Clonality Monoclonal
Isotype IgG1

Calculated MW 61.4kDa KDa

**Description** 

The protein encoded by this gene belongs to the cyclic nucleotide phosphodiesterase (PDE) family, and PDE1 subfamily. Members of the PDE1 family are calmodulin-dependent PDEs that are stimulated by a calcium-calmodulin complex. This PDE has dual-specificity for the second messengers, cAMP and cGMP, with a preference for cGMP as a substrate. cAMP and cGMP function as key regulators of many important physiological processes. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.

#### **Immunogen**

Purified recombinant fragment of human PDE1B (AA: 370-536) expressed in E. Coli.

#### Formulation

Purified antibody in PBS with 0.05% sodium azide

#### **PDE1B Antibody - Additional Information**

**Gene ID** 5153

#### **Other Names**

Calcium/calmodulin-dependent 3', 5'-cyclic nucleotide phosphodiesterase 1B, Cam-PDE 1B, 3.1.4.17, 63 kDa Cam-PDE, PDE1B, PDE1B1, PDE51B

## **Dilution**

WB~~1/500 - 1/2000 FC~~1/200 - 1/400 E~~1/10000

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

PDE1B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **PDE1B Antibody - Protein Information**



## Name PDE1B (HGNC:8775)

# Synonyms PDES1B

#### **Function**

Cyclic nucleotide phosphodiesterase with a dual specificity for the second messengers cAMP and cGMP, which are key regulators of many important physiological processes (PubMed:<a href="http://www.uniprot.org/citations/15260978" target="\_blank">15260978</a>, PubMed:<a href="http://www.uniprot.org/citations/8855339" target="\_blank">8855339</a>, PubMed:<a href="http://www.uniprot.org/citations/9419816" target="\_blank">9419816</a>). Has a preference for cGMP as a substrate (PubMed:<a href="http://www.uniprot.org/citations/9419816" target="\_blank">9419816</a>).


#### **Cellular Location**

Cytoplasm, cytosol.

## **PDE1B Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture








Figure 1: Western blot analysis using PDE1B mAb against human PDE1B recombinant protein. (Expected MW is 44.4 kDa)




Figure 2: Western blot analysis using PDE1B mAb against HEK293 (1) and PDE1B (AA: 370-536)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using PDE1B mouse mAb against A549 (1), SK-MES-1 (2), PC-12 (4),3T3L1 (5) cell lysate and Mouse brain (3) tissue lysate.






Figure 4: Flow cytometric analysis of A549 cells using PDE1B mouse mAb (green) and negative control (red).

# **PDE1B Antibody - References**

1.J Biol Chem. 2007 Nov 9;282(45):32749-57. 2.Proc Natl Acad Sci U S A. 2005 Jan 11;102(2):497-502.