

CK5 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2288a**Specification****CK5 Antibody - Product Information**

Application	WB, IHC, FC, E
Primary Accession	P13647
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	62.4kDa KDa

Description

The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.

Immunogen

Purified recombinant fragment of human CK5 (AA: 316-590) expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

CK5 Antibody - Additional Information

Gene ID 3852

Other Names

Keratin, type II cytoskeletal 5, 58 kDa cytokeratin, Cytokeratin-5, CK-5, Keratin-5, K5, Type-II keratin Kb5, KRT5

Dilution

WB~~1/500 - 1/2000

IHC~~1/200 - 1/1000

FC~~1/200 - 1/400

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CK5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CK5 Antibody - Protein Information

Name KRT5

Function

Required for the formation of keratin intermediate filaments in the basal epidermis and maintenance of the skin barrier in response to mechanical stress (By similarity). Regulates the recruitment of Langerhans cells to the epidermis, potentially by modulation of the abundance of macrophage chemotactic cytokines, macrophage inflammatory cytokines and CTNND1 localization in keratinocytes (By similarity).

Cellular Location

Cytoplasm.


Tissue Location

Expressed in corneal epithelium (at protein level) (PubMed:26758872). Expressed in keratinocytes (at protein level) (PubMed:20128788, PubMed:31302245).

CK5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CK5 mAb against human CK5 recombinant protein. (Expected MW is 47.8 kDa)




Figure 2: Western blot analysis using CK5 mouse mAb against A431 (1) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using CK5 mouse mAb (green) and negative control (red).




Figure 4: Immunohistochemical analysis of paraffin-embedded prostate tissues using CK5 mouse mAb with DAB staining.




Figure 5: Immunohistochemical analysis of paraffin-embedded esophageal cancer tissues using CK5 mouse mAb with DAB staining.

CK5 Antibody - References

1. Am J Surg Pathol. 2009 Nov;33(11):1615-23.
2. J Dermatol. 2009 Aug;36(8):447-52.