

Mouse Monoclonal Antibody to XRN2
Purified Mouse Monoclonal Antibody
Catalog # AO2346a**Specification**

Mouse Monoclonal Antibody to XRN2 - Product Information

Application	WB, IHC, E
Primary Accession	O9H0D6
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	108.5kDa KDa

Description

This gene encodes a 5'-3' exonuclease that promotes transcription termination at cotranscriptional cleavage sites. Alternative splicing results in multiple transcript variants encoding different isoforms.;

Immunogen

Purified recombinant fragment of human XRN2 (AA: 398-547) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

Application Note

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000;

Mouse Monoclonal Antibody to XRN2 - Additional Information

Gene ID 22803

Dilution

WB~~1:1000
IHC~~1:100~500
E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Monoclonal Antibody to XRN2 is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Monoclonal Antibody to XRN2 - Protein Information

Name XRN2

Function

Possesses 5'→3' exoribonuclease activity (By similarity). May promote the termination of transcription by RNA polymerase II. During transcription termination, cleavage at the polyadenylation site liberates a 5' fragment which is subsequently processed to form the mature mRNA and a 3' fragment which remains attached to the elongating polymerase. The processive degradation of this 3' fragment by this protein may promote termination of transcription. Binds to RNA polymerase II (RNAP II) transcription termination R-loops formed by G-rich pause sites (PubMed: [21700224](http://www.uniprot.org/citations/21700224)).

Cellular Location

Nucleus, nucleolus.

Tissue Location


Expressed in the spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocytes, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas Isoform 2 is expressed predominantly in peripheral blood leukocytes

Mouse Monoclonal Antibody to XRN2 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Mouse Monoclonal Antibody to XRN2 - Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using XRN2 mAb against human XRN2 (AA: 398-547) recombinant protein. (Expected MW is 43.1 kDa)




Western blot analysis using XRN2 mAb against HEK293 (1) and XRN2 (AA: 398-547)-hlgGfC transfected HEK293 (2) cell lysate.



Western blot analysis using XRN2 mouse mAb against Raw264.7 (1), HEK293 (2), NTERA-2 (3), LNCap (4), HepG2 (5), HEK293 (6), and Hela (7) cell lysate.



Immunohistochemical analysis of paraffin-embedded colon cancer tissues using XRN2 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using XRN2 mouse mAb with DAB staining.

Mouse Monoclonal Antibody to XRN2 - References

1.EMBO J. 2012 May 30;31(11):2566-78. ; 2.DNA Seq. 2005 Apr;16(2):143-6.;