

Mouse Monoclonal Antibody to POMC

Purified Mouse Monoclonal Antibody Catalog # AO2377a

Specification

Mouse Monoclonal Antibody to POMC - Product Information

Application WB, ICC, E
Primary Accession P01189
Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype Mouse IgG1
Calculated MW 29.4kDa KDa

Description

This gene encodes a preproprotein that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the preproprotein and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. The antimicrobial melanotropin alpha peptide exhibits antibacterial and antifungal activity. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described.;

Immunogen

Purified recombinant fragment of human POMC (AA: 1-150) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

Application Note

ELISA: 1/10000; WB: 1/500 - 1/2000; ICC: 1/200 - 1/1000;

Mouse Monoclonal Antibody to POMC - Additional Information

Gene ID 5443

Other Names

LPH; MSH; NPP; POC; ACTH; CLIP

Dilution
WB~~1:1000
ICC~~N/A



E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Monoclonal Antibody to POMC is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Monoclonal Antibody to POMC - Protein Information

Name POMC

Function

[Corticotropin]: Stimulates the adrenal glands to release cortisol. [Melanocyte-stimulating hormone beta]: Increases the pigmentation of skin by increasing melanin production in melanocytes. [Met-enkephalin]: Endogenous opiate.

Cellular Location

Secreted {ECO:0000250|UniProtKB:P01193}. Note=Melanocyte-stimulating hormone alpha and beta-endorphin are stored in separate granules in hypothalamic POMC neurons, suggesting that secretion may be under the control of different regulatory mechanisms {ECO:0000250|UniProtKB:P01193}

Tissue Location

ACTH and MSH are produced by the pituitary gland.


Mouse Monoclonal Antibody to POMC - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Monoclonal Antibody to POMC - Images






Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Western blot analysis using POMC mAb against human POMC (AA: 1-150) recombinant protein. (Expected MW is 41.9 kDa)



Western blot analysis using POMC mAb against HEK293 (1) and POMC (AA: 1-150)-hlgGFc transfected HEK293 (2) cell lysate.





Immunofluorescence analysis of Hela cells using POMC mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher

Mouse Monoclonal Antibody to POMC - References

1.Tumour Biol. 2015 Mar;36(3):1811-7.; 2.J Neurosci. 2013 Feb 20;33(8):3624-32.;