

Mouse Monoclonal Antibody to PTPN14

Purified Mouse Monoclonal Antibody Catalog # AO2383a

Specification

Mouse Monoclonal Antibody to PTPN14 - Product Information

Application E, WB, FC, ICC, IHC

Primary Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

Q15678
Human
Mouse
Mouse
Monoclonal
Mouse IgG1
135.3kDa KDa

Description

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an N-terminal noncatalytic domain similar to that of band 4.1 superfamily cytoskeleton-associated proteins, which suggested the membrane or cytoskeleton localization of this protein. It appears to regulate lymphatic development in mammals, and a loss of function mutation has been found in a kindred with a lymphedema-choanal atresia.;

Immunogen

Purified recombinant fragment of human PTPN14 (AA: 896-1169) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

Application Note

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 - 1/400

Mouse Monoclonal Antibody to PTPN14 - Additional Information

Gene ID 5784

Other Names PEZ; PTP36

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Monoclonal Antibody to PTPN14 is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Monoclonal Antibody to PTPN14 - Protein Information



Name PTPN14

Synonyms PEZ, PTPD2

Function

Protein tyrosine phosphatase which may play a role in the regulation of lymphangiogenesis, cell-cell adhesion, cell-matrix adhesion, cell migration, cell growth and also regulates TGF-beta gene expression, thereby modulating epithelial-mesenchymal transition. Mediates beta-catenin dephosphorylation at adhesion junctions. Acts as a negative regulator of the oncogenic property of YAP, a downstream target of the hippo pathway, in a cell density-dependent manner. May function as a tumor suppressor.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus. Note=Translocation into the nucleus is associated with induction of cell proliferation. Partially colocalized with actin filaments at the plasma membrane


Tissue Location Ubiquitous.

Mouse Monoclonal Antibody to PTPN14 - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Monoclonal Antibody to PTPN14 - Images




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)






Western blot analysis using PTPN14 mAb against human PTPN14 (AA: 896-1169) recombinant protein. (Expected MW is 57.5 kDa)




Western blot analysis using PTPN14 mAb against HEK293 (1) and PTPN14 (AA: 896-1169)-hlgGFc transfected HEK293 (2) cell lysate.




Flow cytometric analysis of Hela cells using PTPN14 mouse mAb (green) and negative control




(red).




Immunofluorescence analysis of Hela cells using PTPN14 mouse mAb. Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin.



Immunofluorescence analysis of Hela cells using PTPN14 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher



Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using PTPN14 mouse





mAb with DAB staining.

Mouse Monoclonal Antibody to PTPN14 - References

1.Oncogene. 2013 Apr 18;32(16):2087-95.; 2.PLoS One. 2013 Apr 16;8(4):e61916.;