

SMARCA1
Purified Mouse Monoclonal Antibody
Catalog # AO2533a**Specification****SMARCA1 - Product Information**

Application	WB, IHC, ICC, E
Primary Accession	P28370
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	122.6kDa KDa

Immunogen

Purified recombinant fragment of human SMARCA1 (AA: 933-1070) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

SMARCA1 - Additional Information**Gene ID 6594****Other Names**

SWI; ISWI; SWI2; SNF2L; SNF2L1; SNF2LB; SNF2LT; hSNF2L; NURF140

Dilution

WB~~ 1/500 - 1/2000
IHC~~ 1/200 - 1/1000
ICC~~N/A
E~~ 1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SMARCA1 is for research use only and not for use in diagnostic or therapeutic procedures.

SMARCA1 - Protein Information**Name** SMARCA1 ([HGNC:11097](#))**Synonyms** SNF2L, SNF2L1**Function**

[Isoform 1]: ATPase that possesses intrinsic ATP-dependent chromatin-remodeling activity

(PubMed:14609955, PubMed:15310751, PubMed:15640247, PubMed:28801535). ATPase activity is substrate- dependent, and is increased when nucleosomes are the substrate, but is also catalytically active when DNA alone is the substrate (PubMed:14609955, PubMed:15310751, PubMed:15640247). Catalytic subunit of ISWI chromatin-remodeling complexes, which form ordered nucleosome arrays on chromatin and facilitate access to DNA during DNA-templated processes such as DNA replication, transcription, and repair (PubMed:14609955, PubMed:15310751, PubMed:15640247, PubMed:28801535). Within the ISWI chromatin-remodeling complexes, slides edge- and center-positioned histone octamers away from their original location on the DNA template (PubMed:28801535). Catalytic activity and histone octamer sliding propensity is regulated and determined by components of the ISWI chromatin-remodeling complexes (PubMed:28801535). The BAZ1A-, BAZ1B-, BAZ2A- and BAZ2B-containing ISWI chromatin-remodeling complexes regulate the spacing of nucleosomes along the chromatin and have the ability to slide mononucleosomes to the center of a DNA template (PubMed:28801535). The CECR2- and RSF1-containing ISWI chromatin- remodeling complexes do not have the ability to slide mononucleosomes to the center of a DNA template (PubMed:28801535). Within the NURF-1 and CERF-1 ISWI chromatin remodeling complexes, nucleosomes are the preferred substrate for its ATPase activity (PubMed:14609955, PubMed:15640247). Within the NURF-1 ISWI chromatin-remodeling complex, binds to the promoters of En1 and En2 to positively regulate their expression and promote brain development (PubMed:14609955). May promote neurite outgrowth (PubMed:14609955). May be involved in the development of luteal cells (PubMed:16740656). Facilitates nucleosome assembly during DNA replication, ensuring replication fork progression and genomic stability by preventing replication stress and nascent DNA gaps (PubMed:39413208).

Cellular Location

Nucleus. Chromosome

Tissue Location

[Isoform 1]: Expressed in lung, breast, kidney, ovary, skeletal muscle and brain.

SMARCA1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SMARCA1 - Images

Figure 1: Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Figure 2: Western blot analysis using SMARCA1 mAb against human SMARCA1 (AA: 933-1070) recombinant protein. (Expected MW is 42.4 kDa)




Figure 3:Western blot analysis using SMARCA1 mAb against HEK293 (1) and SMARCA1 (AA: 933-1070)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 4:Western blot analysis using SMARCA1 mouse mAb against SW620 (1) and HT-29 (2) cell lysate.




Figure 5:Flow cytometric analysis of NIH/3T3 cells using SMARCA1 mouse mAb (green) and

negative control (red).




Figure 6: Immunohistochemical analysis of paraffin-embedded esophageal cancer tissues using SMARCA1 mouse mAb with DAB staining.




Figure 7: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using SMARCA1 mouse mAb with DAB staining.

SMARCA1 - References

- 1.Yonsei Med J. 2013 May 1;54(3):772-7.2.BMC Med Genet. 2008 Feb 26;9:11.